

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C12N 9/00	A2	(11) International Publication Number: WO 00/32757
		(43) International Publication Date: 8 June 2000 (08.06.00)

(21) International Application Number: PCT/US99/28588
(22) International Filing Date: 2 December 1999 (02.12.99)
(30) Priority Data:
60/110,781 3 December 1998 (03.12.98) US
(71) Applicant (for all designated States except US): E.I. DU PONT
DE NEMOURS AND COMPANY [US/US]; 1007 Market
Street, Wilmington, DE 19898 (US).
(72) Inventors; and
(75) Inventors/Applicants (for US only): CAHOON, Rebecca, E.
[US/US]; 2331 West 18th Street, Wilmington, DE 19806
(US). COUGHLAN, Sean, J. [US/US]; 108 Stratton Drive,
Hockessin, DE 19707 (US). MIAO, Guo-Hua [US/US];
202 Cherry Blossom Place, Hockessin, DE 19707 (US).
RAFALSKI, J., Antoni [US/US]; 2028 Longcome Drive,
Wilmington, DE 19810 (US).
(74) Agent: FEULNER, Gregory, J.; E.I. du Pont de Nemours and
Company, Legal Patent Records Center, 1007 Market Street,
Wilmington, DE 19898 (US).

(81) Designated States: AE, AL, AU, BA, BB, BG, BR, CA, CN,
CR, CU, CZ, DM, EE, GD, GE, HR, HU, ID, IL, IN, IS,
JP, KP, KR, LC, LK, LR, LT, LV, MG, MK, MN, MX,
NO, NZ, PL, RO, SG, SI, SK, SL, TR, TT, UA, US, UZ,
VN, YU, ZA, ARIPO patent (GH, GM, KE, LS, MW, SD,
SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG,
KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY,
DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT,
SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW,
ML, MR, NE, SN, TD, TG).

Published

Without international search report and to be republished
upon receipt of that report.

(54) Title: PLANT VITAMIN E BIOSYNTHETIC ENZYMES

(57) Abstract

This invention relates to an isolated nucleic acid fragment encoding a vitamin E biosynthetic enzyme. The invention also relates to the construction of a chimeric gene encoding all or a portion of the vitamin E biosynthetic enzyme, in sense or antisense orientation, wherein expression of the chimeric gene results in production of altered levels of the vitamin E biosynthetic enzyme in a transformed host cell.

SEQ ID NO:39 HYVVRPKNAL-----FLATCYTS
SEQ ID NO:40 KATLAAPSLTSLP-----TKTSS-FCSSLLFRSPSSSSYS-NTTTRNV
SEQ ID NO:41 HAN-----SKPA-PLTPKHLGAAFPFPLGHAAPVPRVPLPLPAR-KLAPGV
SEQ ID NO:42 HATVRI-----PTICRIHTTSGSPRTTARIVGSPNAPTRADA-ASSGCEI
SEQ ID NO:43 HATVRI-----PTICRIHTTSGSPRTTARIVGSPNAPTRADA-ASSGCEI
SEQ ID NO:44 HANAALLCSGDS-GLANCSTHRAHSP-VNHRALRAVVEL-----B-PLASTA-
SEQ ID NO:45 AR-WQPTGALAPLPLACTSRKLCAS3PRAGLCUWHRHRS33HTLLAVRANPT
SEQ ID NO:46 PRR-----GR-----SL-----60
1
SEQ ID NO:39 LTHASATIASADLYEKIKNFYDSSCLNEHDMCHHNGCYGPGTIRL-----DRDQAI
SEQ ID NO:40 AVAAATSTEA-LKGIARTFMTSCLEKINCDHNGCTFQDESVOLESCKEAKAI
SEQ ID NO:41 VDQGGCDAAAPKLECIAGLYDESSGLWETHECHNHGFTDSCGAASHSD-BKRAAI
SEQ ID NO:42 VLQKFAKDEKHLGKCIATFYKESGLKHEHNDHNGFTDSDTVSLSD-BKRAAI
SEQ ID NO:43 VLQKFAKDEKHLGKCIATFYKESGLKHEHNDHNGFTDSDTVSLSD-BKRAAI
SEQ ID NO:44 VLQKFAKDEKHLGKCIATFYKESGLKHEHNDHNGFTDSDTVSLSD-BKRAAI
SEQ ID NO:45 VLQKFAKDEKHLGKCIATFYKESGLKHEHNDHNGFTDSDTVSLSD-BKRAAI
SEQ ID NO:46 VLQKFAKDEKHLGKCIATFYKESGLKHEHNDHNGFTDSDTVSLSD-BKRAAI
120
SEQ ID NO:39 DLTKELAAVAVQKSA-----KPKILGQGGIIGSSLYLAQGHVNGHSLFPQVRA
SEQ ID NO:40 RKIKELAFAGVYDEE-EKRIKVVWGGIIGSSRYLAKFGACIGITLSPVQAKRA
SEQ ID NO:41 RHIEZALATAAVP-DDPTHTPTIYVGGCIGSSRYLAKFGACIGITLSPVQAKRA
SEQ ID NO:42 RHIOELAFASV-EELASRWPHIVDGGCIGSSRYLAKFGACIGITLSPVQAKRA
SEQ ID NO:43 RHIOELAFASV-EELASRWPHIVDGGCIGSSRYLAKFGACIGITLSPVQAKRA
SEQ ID NO:44 RHIEZALATAAVP-DDPTHTPTIYVGGCIGSSRYLAKFGACIGITLSPVQAKRA
SEQ ID NO:45 RHIEZALATAAVP-DDPTHTPTIYVGGCIGSSRYLAKFGACIGITLSPVQAKRA
SEQ ID NO:46 RHIEZALATAAVP-DDPTHTPTIYVGGCIGSSRYLAKFGACIGITLSPVQAKRA
180
SEQ ID NO:39 GERARALGCTCTQTVAMALDLPFASDFQVNSLESGENHPKQAFLOEAVRLKPGG
SEQ ID NO:40 HLAALAGGLSKASITQVADALQDFPQCDLTVWNSLESGENHPKQAFLOEAVRLKPGG
SEQ ID NO:41 HLAALAGGLSKASITQVADALQDFPQCDLTVWNSLESGENHPKQAFLOEAVRLKPGG
SEQ ID NO:42 HLAALAGGLSKASITQVADALQDFPQCDLTVWNSLESGENHPKQAFLOEAVRLKPGG
SEQ ID NO:43 HLAALAGGLSKASITQVADALQDFPQCDLTVWNSLESGENHPKQAFLOEAVRLKPGG
SEQ ID NO:44 HLAALAGGLSKASITQVADALQDFPQCDLTVWNSLESGENHPKQAFLOEAVRLKPGG
SEQ ID NO:45 HLAALAGGLSKASITQVADALQDFPQCDLTVWNSLESGENHPKQAFLOEAVRLKPGG
SEQ ID NO:46 HLAALAGGLSKASITQVADALQDFPQCDLTVWNSLESGENHPKQAFLOEAVRLKPGG
240
SEQ ID NO:39 RLILATWCHRFIDPGHPLTADERRALQAIYDVTCLFVYSLPQTEATACQCGEIKTA
SEQ ID NO:40 RLILATWCHRFIDPGHPLTADERRALQAIYDVTCLFVYSLPQTEATACQCGEIKTA
SEQ ID NO:41 RLILATWCHRFIDPGHPLTADERRALQAIYDVTCLFVYSLPQTEATACQCGEIKTA
SEQ ID NO:42 RLILATWCHRFIDPGHPLTADERRALQAIYDVTCLFVYSLPQTEATACQCGEIKTA
SEQ ID NO:43 RLILATWCHRFIDPGHPLTADERRALQAIYDVTCLFVYSLPQTEATACQCGEIKTA
SEQ ID NO:44 RLILATWCHRFIDPGHPLTADERRALQAIYDVTCLFVYSLPQTEATACQCGEIKTA
SEQ ID NO:45 RLILATWCHRFIDPGHPLTADERRALQAIYDVTCLFVYSLPQTEATACQCGEIKTA
SEQ ID NO:46 RLILATWCHRFIDPGHPLTADERRALQAIYDVTCLFVYSLPQTEATACQCGEIKTA
300
SEQ ID NO:39 QNSVAVFPNHRVIESATQRYVLAQGGPRIIHAALCLAHNGVYKGLVRFGLATCI
SEQ ID NO:40 QNSVAVFPNHRVIESATQRYVLAQGGPRIIHAALCLAHNGVYKGLVRFGLATCI
SEQ ID NO:41 QNSVAVFPNHRVIESATQRYVLAQGGPRIIHAALCLAHNGVYKGLVRFGLATCI
SEQ ID NO:42 QNSVAVFPNHRVIESATQRYVLAQGGPRIIHAALCLAHNGVYKGLVRFGLATCI
SEQ ID NO:43 QNSVAVFPNHRVIESATQRYVLAQGGPRIIHAALCLAHNGVYKGLVRFGLATCI
SEQ ID NO:44 QNSVAVFPNHRVIESATQRYVLAQGGPRIIHAALCLAHNGVYKGLVRFGLATCI
SEQ ID NO:45 QNSVAVFPNHRVIESATQRYVLAQGGPRIIHAALCLAHNGVYKGLVRFGLATCI
SEQ ID NO:46 QNSVAVFPNHRVIESATQRYVLAQGGPRIIHAALCLAHNGVYKGLVRFGLATCI
360
SEQ ID NO:39 EPLV-----
SEQ ID NO:40 EPLV-----
SEQ ID NO:41 EPLV-----
SEQ ID NO:42 EPLV-----
SEQ ID NO:43 EPLV-----
SEQ ID NO:44 EPLV-----
SEQ ID NO:45 EPLV-----
SEQ ID NO:46 EPLV-----
380
SEQ ID NO:39 EPLV-----
SEQ ID NO:40 EPLV-----
SEQ ID NO:41 EPLV-----
SEQ ID NO:42 EPLV-----
SEQ ID NO:43 EPLV-----
SEQ ID NO:44 EPLV-----
SEQ ID NO:45 EPLV-----
SEQ ID NO:46 EPLV-----
400
SEQ ID NO:39 EPLV-----
SEQ ID NO:40 EPLV-----
SEQ ID NO:41 EPLV-----
SEQ ID NO:42 EPLV-----
SEQ ID NO:43 EPLV-----
SEQ ID NO:44 EPLV-----
SEQ ID NO:45 EPLV-----
SEQ ID NO:46 EPLV-----
420
SEQ ID NO:39 EPLV-----
SEQ ID NO:40 EPLV-----
SEQ ID NO:41 EPLV-----
SEQ ID NO:42 EPLV-----
SEQ ID NO:43 EPLV-----
SEQ ID NO:44 EPLV-----
SEQ ID NO:45 EPLV-----
SEQ ID NO:46 EPLV-----
440
SEQ ID NO:39 EPLV-----
SEQ ID NO:40 EPLV-----
SEQ ID NO:41 EPLV-----
SEQ ID NO:42 EPLV-----
SEQ ID NO:43 EPLV-----
SEQ ID NO:44 EPLV-----
SEQ ID NO:45 EPLV-----
SEQ ID NO:46 EPLV-----
460
SEQ ID NO:39 EPLV-----
SEQ ID NO:40 EPLV-----
SEQ ID NO:41 EPLV-----
SEQ ID NO:42 EPLV-----
SEQ ID NO:43 EPLV-----
SEQ ID NO:44 EPLV-----
SEQ ID NO:45 EPLV-----
SEQ ID NO:46 EPLV-----
480
SEQ ID NO:39 EPLV-----
SEQ ID NO:40 EPLV-----
SEQ ID NO:41 EPLV-----
SEQ ID NO:42 EPLV-----
SEQ ID NO:43 EPLV-----
SEQ ID NO:44 EPLV-----
SEQ ID NO:45 EPLV-----
SEQ ID NO:46 EPLV-----
500
SEQ ID NO:39 EPLV-----
SEQ ID NO:40 EPLV-----
SEQ ID NO:41 EPLV-----
SEQ ID NO:42 EPLV-----
SEQ ID NO:43 EPLV-----
SEQ ID NO:44 EPLV-----
SEQ ID NO:45 EPLV-----
SEQ ID NO:46 EPLV-----
520
SEQ ID NO:39 EPLV-----
SEQ ID NO:40 EPLV-----
SEQ ID NO:41 EPLV-----
SEQ ID NO:42 EPLV-----
SEQ ID NO:43 EPLV-----
SEQ ID NO:44 EPLV-----
SEQ ID NO:45 EPLV-----
SEQ ID NO:46 EPLV-----
540
SEQ ID NO:39 EPLV-----
SEQ ID NO:40 EPLV-----
SEQ ID NO:41 EPLV-----
SEQ ID NO:42 EPLV-----
SEQ ID NO:43 EPLV-----
SEQ ID NO:44 EPLV-----
SEQ ID NO:45 EPLV-----
SEQ ID NO:46 EPLV-----
560
SEQ ID NO:39 EPLV-----
SEQ ID NO:40 EPLV-----
SEQ ID NO:41 EPLV-----
SEQ ID NO:42 EPLV-----
SEQ ID NO:43 EPLV-----
SEQ ID NO:44 EPLV-----
SEQ ID NO:45 EPLV-----
SEQ ID NO:46 EPLV-----
580
SEQ ID NO:39 EPLV-----
SEQ ID NO:40 EPLV-----
SEQ ID NO:41 EPLV-----
SEQ ID NO:42 EPLV-----
SEQ ID NO:43 EPLV-----
SEQ ID NO:44 EPLV-----
SEQ ID NO:45 EPLV-----
SEQ ID NO:46 EPLV-----
600
SEQ ID NO:39 EPLV-----
SEQ ID NO:40 EPLV-----
SEQ ID NO:41 EPLV-----
SEQ ID NO:42 EPLV-----
SEQ ID NO:43 EPLV-----
SEQ ID NO:44 EPLV-----
SEQ ID NO:45 EPLV-----
SEQ ID NO:46 EPLV-----
620
SEQ ID NO:39 EPLV-----
SEQ ID NO:40 EPLV-----
SEQ ID NO:41 EPLV-----
SEQ ID NO:42 EPLV-----
SEQ ID NO:43 EPLV-----
SEQ ID NO:44 EPLV-----
SEQ ID NO:45 EPLV-----
SEQ ID NO:46 EPLV-----
640
SEQ ID NO:39 EPLV-----
SEQ ID NO:40 EPLV-----
SEQ ID NO:41 EPLV-----
SEQ ID NO:42 EPLV-----
SEQ ID NO:43 EPLV-----
SEQ ID NO:44 EPLV-----
SEQ ID NO:45 EPLV-----
SEQ ID NO:46 EPLV-----
660
SEQ ID NO:39 EPLV-----
SEQ ID NO:40 EPLV-----
SEQ ID NO:41 EPLV-----
SEQ ID NO:42 EPLV-----
SEQ ID NO:43 EPLV-----
SEQ ID NO:44 EPLV-----
SEQ ID NO:45 EPLV-----
SEQ ID NO:46 EPLV-----
680
SEQ ID NO:39 EPLV-----
SEQ ID NO:40 EPLV-----
SEQ ID NO:41 EPLV-----
SEQ ID NO:42 EPLV-----
SEQ ID NO:43 EPLV-----
SEQ ID NO:44 EPLV-----
SEQ ID NO:45 EPLV-----
SEQ ID NO:46 EPLV-----
700
SEQ ID NO:39 EPLV-----
SEQ ID NO:40 EPLV-----
SEQ ID NO:41 EPLV-----
SEQ ID NO:42 EPLV-----
SEQ ID NO:43 EPLV-----
SEQ ID NO:44 EPLV-----
SEQ ID NO:45 EPLV-----
SEQ ID NO:46 EPLV-----
720
SEQ ID NO:39 EPLV-----
SEQ ID NO:40 EPLV-----
SEQ ID NO:41 EPLV-----
SEQ ID NO:42 EPLV-----
SEQ ID NO:43 EPLV-----
SEQ ID NO:44 EPLV-----
SEQ ID NO:45 EPLV-----
SEQ ID NO:46 EPLV-----
740
SEQ ID NO:39 EPLV-----
SEQ ID NO:40 EPLV-----
SEQ ID NO:41 EPLV-----
SEQ ID NO:42 EPLV-----
SEQ ID NO:43 EPLV-----
SEQ ID NO:44 EPLV-----
SEQ ID NO:45 EPLV-----
SEQ ID NO:46 EPLV-----
760
SEQ ID NO:39 EPLV-----
SEQ ID NO:40 EPLV-----
SEQ ID NO:41 EPLV-----
SEQ ID NO:42 EPLV-----
SEQ ID NO:43 EPLV-----
SEQ ID NO:44 EPLV-----
SEQ ID NO:45 EPLV-----
SEQ ID NO:46 EPLV-----
780
SEQ ID NO:39 EPLV-----
SEQ ID NO:40 EPLV-----
SEQ ID NO:41 EPLV-----
SEQ ID NO:42 EPLV-----
SEQ ID NO:43 EPLV-----
SEQ ID NO:44 EPLV-----
SEQ ID NO:45 EPLV-----
SEQ ID NO:46 EPLV-----
800
SEQ ID NO:39 EPLV-----
SEQ ID NO:40 EPLV-----
SEQ ID NO:41 EPLV-----
SEQ ID NO:42 EPLV-----
SEQ ID NO:43 EPLV-----
SEQ ID NO:44 EPLV-----
SEQ ID NO:45 EPLV-----
SEQ ID NO:46 EPLV-----
820
SEQ ID NO:39 EPLV-----
SEQ ID NO:40 EPLV-----
SEQ ID NO:41 EPLV-----
SEQ ID NO:42 EPLV-----
SEQ ID NO:43 EPLV-----
SEQ ID NO:44 EPLV-----
SEQ ID NO:45 EPLV-----
SEQ ID NO:46 EPLV-----
840
SEQ ID NO:39 EPLV-----
SEQ ID NO:40 EPLV-----
SEQ ID NO:41 EPLV-----
SEQ ID NO:42 EPLV-----
SEQ ID NO:43 EPLV-----
SEQ ID NO:44 EPLV-----
SEQ ID NO:45 EPLV-----
SEQ ID NO:46 EPLV-----
860
SEQ ID NO:39 EPLV-----
SEQ ID NO:40 EPLV-----
SEQ ID NO:41 EPLV-----
SEQ ID NO:42 EPLV-----
SEQ ID NO:43 EPLV-----
SEQ ID NO:44 EPLV-----
SEQ ID NO:45 EPLV-----
SEQ ID NO:46 EPLV-----
880
SEQ ID NO:39 EPLV-----
SEQ ID NO:40 EPLV-----
SEQ ID NO:41 EPLV-----
SEQ ID NO:42 EPLV-----
SEQ ID NO:43 EPLV-----
SEQ ID NO:44 EPLV-----
SEQ ID NO:45 EPLV-----
SEQ ID NO:46 EPLV-----
900
SEQ ID NO:39 EPLV-----
SEQ ID NO:40 EPLV-----
SEQ ID NO:41 EPLV-----
SEQ ID NO:42 EPLV-----
SEQ ID NO:43 EPLV-----
SEQ ID NO:44 EPLV-----
SEQ ID NO:45 EPLV-----
SEQ ID NO:46 EPLV-----
920
SEQ ID NO:39 EPLV-----
SEQ ID NO:40 EPLV-----
SEQ ID NO:41 EPLV-----
SEQ ID NO:42 EPLV-----
SEQ ID NO:43 EPLV-----
SEQ ID NO:44 EPLV-----
SEQ ID NO:45 EPLV-----
SEQ ID NO:46 EPLV-----
940
SEQ ID NO:39 EPLV-----
SEQ ID NO:40 EPLV-----
SEQ ID NO:41 EPLV-----
SEQ ID NO:42 EPLV-----
SEQ ID NO:43 EPLV-----
SEQ ID NO:44 EPLV-----
SEQ ID NO:45 EPLV-----
SEQ ID NO:46 EPLV-----
960
SEQ ID NO:39 EPLV-----
SEQ ID NO:40 EPLV-----
SEQ ID NO:41 EPLV-----
SEQ ID NO:42 EPLV-----
SEQ ID NO:43 EPLV-----
SEQ ID NO:44 EPLV-----
SEQ ID NO:45 EPLV-----
SEQ ID NO:46 EPLV-----
980
SEQ ID NO:39 EPLV-----
SEQ ID NO:40 EPLV-----
SEQ ID NO:41 EPLV-----
SEQ ID NO:42 EPLV-----
SEQ ID NO:43 EPLV-----
SEQ ID NO:44 EPLV-----
SEQ ID NO:45 EPLV-----
SEQ ID NO:46 EPLV-----
1000
SEQ ID NO:39 EPLV-----
SEQ ID NO:40 EPLV-----
SEQ ID NO:41 EPLV-----
SEQ ID NO:42 EPLV-----
SEQ ID NO:43 EPLV-----
SEQ ID NO:44 EPLV-----
SEQ ID NO:45 EPLV-----
SEQ ID NO:46 EPLV-----

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav	TM	Turkmenistan
BF	Burkina Faso	GR	Greece		Republic of Macedonia	TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's	NZ	New Zealand		
CM	Cameroon		Republic of Korea	PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

TITLE

PLANT VITAMIN E BIOSYNTHETIC ENZYMES

This application claims the benefit of U.S. Provisional Application No. 60/110,781, filed December 3, 1998.

FIELD OF THE INVENTION

This invention is in the field of plant molecular biology. More specifically, this invention pertains to nucleic acid fragments encoding vitamin E biosynthetic enzymes in plants and seeds.

BACKGROUND OF THE INVENTION

10 Vitamin E (alpha-tocopherol) is an essential element in the mammalian diet since mammals can not synthesize plastoquinones or tocopherols. The first step in the formation of plastoquinones and tocopherols in plants is the formation of homogениstate from 4-hydroxyphenylpyruvate, a reaction catalyzed by 4-hydroxyphenylpyruvate dioxygenase (EC 1.13.11.27). Genetic mutants of this enzyme in *Arabidopsis* are deficient in both
15 carotenoids and tocopherols (Norris et al. (1995) *Plant Cell* 7:2139-2149). Carotenoids (xanthophylls) in corn endosperm are valuable coloring agents in poultry feeds; tocopherols are antioxidants which may be important in oil stability and human health. Generally present at low levels in plant tissues, 4-hydroxyphenylpyruvate dioxygenase catalyzes a complex, irreversible reaction. Only recently has 4-hydroxyphenylpyruvate dioxygenase
20 been purified to homogeneity from a plant source (Garcia et al. (1997 *Biochem. J.* 325:761-769). In plants, cDNAs encoding 4-hydroxyphenylpyruvate dioxygenase have been identified in carrots, barley and *Arabidopsis thaliana* with at least two different variants existing in this last plant. 4-Hydroxyphenylpyruvate dioxygenase is a known herbicide target (Mayonado et al. (1989) *Pestic. Biochem. Physiol.* 35:138-145; Schultz et al. (1993)
25 *FEBS lett.* 318:162-166; Secor (1994) *Plant Phys.* 106:1429-1433). Even though plastoquinones and tocopherols are not synthesized by mammals and bacteria 4-hydroxyphenylpyruvate dioxygenase activity is present, often at high levels, and involved in phenylalanine and tyrosine degradation. Among others, the cDNAs encoding 4-hydroxyphenylpyruvate dioxygenase have been identified in *Mycosphaerella graminicola*,
30 mice and *Coccidioides immitis*.

Plants synthesize alpha, beta, gamma and delta tocopherols of which alpha tocopherol (vitamin E) has the highest value for human nutrition. In soybean 7% of the tocopherols are vitamin E. Gamma tocopherol methyltransferase catalyzes the final step in vitamin E
35 synthesis and has been purified to homogeneity from pepper, marigold, *Euglena* and spinach. The gene encoding gamma tocopherol methyltransferase from *Synechocystis* was identified by mutating an open reading frame encoding a methyltransferase located in the operon containing the 4-hydroxyphenylpyruvate dioxygenase gene in the *Synechocystis*

genomic sequence (Shintani and Della Penna (1998) Abstract for the American Society of Plant Physiologists meeting in Madison, WI).

Since mammals can not synthesize tocopherols, the enzymes described here may be used for the discovery of new herbicides.

5

SUMMARY OF THE INVENTION

The present invention relates to isolated polynucleotides comprising a nucleotide sequence encoding a polypeptide of at least 160 amino acids that has at least 80% identity based on the Clustal method of alignment when compared to a polypeptide selected from the group consisting of a corn gamma-tocopherol methyltransferase polypeptide of SEQ ID
10 NOs:2 and 22, a rice gamma-tocopherol methyltransferase polypeptide of SEQ ID NOs:4, 6, 24, and 26, a soybean gamma-tocopherol methyltransferase polypeptide of SEQ ID NOs:8 and 28, and a wheat gamma-tocopherol methyltransferase polypeptide of SEQ ID NOs:10 and 30. The present invention also relates to an isolated polynucleotide comprising the complement of the nucleotide sequences described above.

15 The present invention relates to isolated polynucleotides comprising a nucleotide sequence encoding a polypeptide of at least 150 amino acids that has at least 95% identity based on the Clustal method of alignment when compared to a polypeptide selected from the group consisting of a catalpa 4-hydroxyphenylpyruvate dioxygenase polypeptide of SEQ ID NO:32, a rice 4-hydroxyphenylpyruvate dioxygenase polypeptide of SEQ ID NOs:12, 14,
20 and 34, a soybean 4-hydroxyphenylpyruvate dioxygenase polypeptide of SEQ ID NOs:16 and 36, a Vernonia 4-hydroxyphenylpyruvate dioxygenase of SEQ ID NO:18, and a wheat 4-hydroxyphenylpyruvate dioxygenase polypeptide of SEQ ID NOs:20 and 38. The present invention also relates to an isolated polynucleotide comprising the complement of the nucleotide sequences described above.

25 It is preferred that the isolated polynucleotides of the claimed invention consist of a nucleic acid sequence selected from the group consisting of SEQ ID NOs:1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, and 37 that codes for the polypeptide selected from the group consisting of SEQ ID NOs:2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, and 38. The present invention also relates to an isolated polynucleotide
30 comprising a nucleotide sequences of at least one of 60 (preferably at least one of 40, most preferably at least one of 30) contiguous nucleotides derived from a nucleotide sequence selected from the group consisting of SEQ ID NOs:1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, and the complement of such nucleotide sequences.

The present invention relates to a chimeric gene comprising an isolated polynucleotide
35 of the present invention operably linked to suitable regulatory sequences.

The present invention relates to an isolated host cell comprising a chimeric gene of the present invention or an isolated polynucleotide of the present invention. The host cell may be eukaryotic, such as a yeast or a plant cell, or prokaryotic, such as a bacterial cell. The

present invention also relates to a virus, preferably a baculovirus, comprising an isolated polynucleotide of the present invention or a chimeric gene of the present invention.

The present invention relates to a process for producing an isolated host cell comprising a chimeric gene of the present invention or an isolated polynucleotide of the present invention, the process comprising either transforming or transfecting an isolated
5 compatible host cell with a chimeric gene or isolated polynucleotide of the present invention.

The present invention relates to a gamma-tocopherol methyltransferase polypeptide of at least 160 amino acids comprising at least 80% homology based on the Clustal method of alignment compared to a polypeptide selected from the group consisting of SEQ ID NOs:2,
10 4, 6, 8, 10, 22, 24, 26, 28, and 30.

The present invention relates to a 4-hydroxyphenylpyruvate dioxygenase polypeptide of at least 150 amino acids comprising at least 95% homology based on the Clustal method of alignment compared to a polypeptide selected from the group consisting of SEQ ID NOs:12, 14, 16, 18, 20, 32, 34, 36, and 38.

15 The present invention relates to a method of selecting an isolated polynucleotide that affects the level of expression of a gamma-tocopherol methyltransferase or a 4-hydroxyphenylpyruvate dioxygenase polypeptide in a host cell, preferably a plant cell, the method comprising the steps of: (a) constructing an isolated polynucleotide of the present invention or an isolated chimeric gene of the present invention; (b) introducing the isolated
20 polynucleotide or the isolated chimeric gene into a host cell; (c) measuring the level a gamma-tocopherol methyltransferase or a 4-hydroxyphenylpyruvate dioxygenase polypeptide in the host cell containing the isolated polynucleotide; and (d) comparing the level of a gamma-tocopherol methyltransferase or a 4-hydroxyphenylpyruvate dioxygenase polypeptide in the host cell containing the isolated polynucleotide with the level of a gamma-
25 tocopherol methyltransferase or a 4-hydroxyphenylpyruvate dioxygenase polypeptide in the host cell that does not contain the isolated polynucleotide.

The present invention relates to a method of obtaining a nucleic acid fragment encoding a substantial portion of a gamma-tocopherol methyltransferase or a 4-hydroxyphenylpyruvate dioxygenase polypeptide gene, preferably a plant gamma-tocopherol
30 methyltransferase or 4-hydroxyphenylpyruvate dioxygenase polypeptide gene, comprising the steps of : synthesizing an oligonucleotide primer comprising a nucleotide sequence of at least one of 60 (preferably at least one of 40, most preferably at least one of 30) contiguous nucleotides derived from a nucleotide sequence selected from the group consisting of SEQ ID NOs:1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, and the
35 complement of such nucleotide sequences; and amplifying a nucleic acid fragment (preferably a cDNA inserted in a cloning vector) using the oligonucleotide primer. The amplified nucleic acid fragment preferably will encode a portion of a gamma-tocopherol methyltransferase or a 4-hydroxyphenylpyruvate dioxygenase amino acid sequence.

The present invention also relates to a method of obtaining a nucleic acid fragment encoding all or a substantial portion of the amino acid sequence encoding a 4-hydroxyphenylpyruvate dioxygenase or a gamma tocopherol methyltransferase polypeptide comprising the steps of: probing a cDNA or genomic library with an isolated polynucleotide of the present invention; identifying a DNA clone that hybridizes with an isolated polynucleotide of the present invention; isolating the identified DNA clone; and sequencing the cDNA or genomic fragment that comprises the isolated DNA clone.

A further embodiment of the instant invention is a method for evaluating at least one compound for its ability to inhibit the activity of a 4-hydroxyphenylpyruvate dioxygenase or a gamma tocopherol methyltransferase, the method comprising the steps of:

- (a) transforming a host cell with a chimeric gene comprising a nucleic acid fragment encoding a 4-hydroxyphenylpyruvate dioxygenase or a gamma tocopherol methyltransferase, operably linked to suitable regulatory sequences;
- (b) growing the transformed host cell under conditions that are suitable for expression of the chimeric gene wherein expression of the chimeric gene results in production of 4-hydroxyphenylpyruvate dioxygenase or gamma tocopherol methyltransferase in the transformed host cell;
- (c) optionally purifying the 4-hydroxyphenylpyruvate dioxygenase or the gamma tocopherol methyltransferase expressed by the transformed host cell;
- (d) treating the 4-hydroxyphenylpyruvate dioxygenase or the gamma tocopherol methyltransferase with a compound to be tested; and
- (e) comparing the activity of the 4-hydroxyphenylpyruvate dioxygenase or the gamma tocopherol methyltransferase that has been treated with a test compound to the activity of an untreated 4-hydroxyphenylpyruvate dioxygenase or gamma tocopherol methyltransferase, thereby selecting compounds with potential for inhibitory activity.

The present invention relates to a composition, such as a hybridization mixture, comprising an isolated polynucleotide of the present invention.

The present invention relates to an isolated polynucleotide of the present invention comprising at least one of 30 contiguous nucleotides derived from a nucleic acid sequence selected from the group consisting of SEQ ID NOs: 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, and 37.

The present invention relates to an expression cassette comprising an isolated polynucleotide of the present invention operably linked to a promoter.

The present invention relates to a method for positive selection of a transformed cell comprising: (a) transforming a host cell with the chimeric gene of the present invention or an expression cassette of the present invention; and (b) growing the transformed host cell, preferably plant cell, such as a monocot or a dicot, under conditions which allow expression of the gamma-tocopherol methyltransferase or a 4-hydroxyphenylpyruvate dioxygenase polynucleotide in an amount sufficient to complement a null mutant to provide a positive selection means.

BRIEF DESCRIPTION OF THE
DRAWINGS AND SEQUENCE DESCRIPTIONS

The invention can be more fully understood from the following detailed description and the accompanying drawings and Sequence Listing which form a part of this application.

5 Figure 1 shows a comparison of the amino acid sequences of the gamma-tocopherol methyltransferase from soybean contig assembled from clones sah1c.pk001.k8, sgc5c.pk001.m23, and sah1c.pk004.g2 (SEQ ID NO:8), corn contig assembled from clones cr1n.pk0179.fl0:fls, cs1.pk0065.f2, ctal1n.pk0031.d2, p0060.coran49r, and p0103.ciaay86r (SEQ ID NO:22), the 3'-end sequence of rice clone rls72.pk0010.g3 (SEQ ID NO:24), the 5'-
10 end sequence of clone rls72.pk0010.g3 (SEQ ID NO:26), soybean clone sah1c.pk004.g2 (SEQ ID NO:28), wheat clone wr1.pk0077.fl:fls (SEQ ID NO:30), *Synechocystis* sp. (NCBI General Identifier No. 1001725; SEQ ID NO:39), and *Arabidopsis thaliana* (NCBI General Identifier No. 4106538; SEQ ID NO:40). Dashes are used by the program to maximize the alignment.

15 Figure 2 shows a comparison of the amino acid sequences of the 4-hydroxyphenyl-pyruvate dioxygenase from catalpa clone ncs.pk0012.g1:fls (SEQ ID NO:32), soybean clone sgc5c.pk001.j9:fls (SEQ ID NO:36), wheat clone wdk4c.pk006.m9:fls (SEQ ID NO:38), *Hordeum vulgare* having NCBI General Identifier No. 3334222 (SEQ ID NO:41), *Daucus carota* having NCBI General Identifier No. 3334219 (SEQ ID NO:42), and *Arabidopsis*
20 *thaliana* having NCBI General Identifier No. 3334223 (SEQ ID NO:43). Dashes are used by the program to maximize the alignment.

Figure 3 depicts the distribution of the percent of alpha- and gamma-tocopherol content of 25 transgenic soybean lines transformed with the gamma tocopherol methyltransferase sequence from clone sah1c.pk001.k8.

25 Table 1 lists the polypeptides that are described herein, the designation of the cDNA clones that comprise the nucleic acid fragments encoding polypeptides representing all or a substantial portion of these polypeptides, and the corresponding identifier (SEQ ID NO:) as used in the attached Sequence Listing. The sequence descriptions and Sequence Listing attached hereto comply with the rules governing nucleotide and/or amino acid sequence
30 disclosures in patent applications as set forth in 37 C.F.R. §1.821-1.825.

TABLE 1
Vitamin E Biosynthetic Enzymes

Protein	Clone Designation	SEQ ID NO:	
		(Nucleotide)	(Amino Acid)
Corn Gamma Tocopherol Methyltransferase	Contig of: cr1n.pk0179.f10 cs1.pk0065.e1 cs1.pk0065.f2 cta1n.pk0031.d2	1	2
Rice Gamma Tocopherol Methyltransferase	rl0n.pk085.e11	3	4
Rice Gamma Tocopherol Methyltransferase	Contig of: rl0n.pk099.d24 rls72.pk0010.g3	5	6
Soybean Gamma Tocopherol Methyltransferase	Contig of: sah1c.pk001.k8 sgc5c.pk001.m23 sah1c.pk004.g2	7	8
Wheat Gamma Tocopherol Methyltransferase	Contig of: wr1.pk0077.f1 wr1.pk177.b11 wle1n.pk0065.h9 wre1n.pk0111.d6	9	10
Rice 4-Hydroxyphenylpyruvate Dioxygenase	rlr12.pk0019.g5	11	12
Rice 4-Hydroxyphenylpyruvate Dioxygenase	rlr12.pk0025.e4	13	14
Soybean 4-Hydroxyphenylpyruvate Dioxygenase	Contig of: sgc5c.pk001.j9 sgs1c.pk002.a8 sfl1.pk126.n15 sgs1c.pk003.o5 sgc4c.pk001.p2	15	16
Vernonia 4-Hydroxyphenylpyruvate Dioxygenase	vs1.pk0015.b2	17	18
Wheat 4-Hydroxyphenylpyruvate Dioxygenase	Contig of: wdk4c.pk006.m9 wlmk1.pk0021.h10 wlmk1.pk0019.e2 wlm0.pk0035.d5 wlmk1.pk0013.g11	19	20
Corn Gamma Tocopherol Methyltransferase	Contig of: cr1n.pk0179.f10: fis cs1.pk0065.f2 cta1n.pk0031.d2 p0060.coran49r p0103.ciaay86r	21	22
Rice Gamma Tocopherol Methyltransferase	rls72.pk0010.g3-3'	23	24

Protein	Clone Designation	SEQ ID NO:	
		(Nucleotide)	(Amino Acid)
Rice Gamma Tocopherol Methyltransferase	rls72.pk0010.g3-5'	25	26
Soybean Gamma Tocopherol Methyltransferase	sah1c.pk004.g2	27	28
Wheat Gamma Tocopherol Methyltransferase	wr1.pk0077.f1: fis	29	30
Catalpa 4-Hydroxyphenyl-pyruvate Dioxygenase	ncs.pk0012.g1: fis	31	32
Rice 4-Hydroxyphenylpyruvate Dioxygenase	rlr12.pk0025.e4: fis	33	34
Soybean 4-Hydroxyphenyl-pyruvate Dioxygenase	sgc5c.pk001.j9: fis	35	36
Wheat 4-Hydroxyphenylpyruvate Dioxygenase	wdk4c.pk006.m9: fis	37	38

The Sequence Listing contains the one letter code for nucleotide sequence characters and the three letter codes for amino acids as defined in conformity with the IUPAC-IUBMB standards described in *Nucleic Acids Res.* 13:3021-3030 (1985) and in the *Biochemical J.* 219 (No. 2):345-373 (1984) which are herein incorporated by reference. The symbols and format used for nucleotide and amino acid sequence data comply with the rules set forth in 37 C.F.R. §1.822.

DETAILED DESCRIPTION OF THE INVENTION

In the context of this disclosure, a number of terms shall be utilized. As used herein, a "polynucleotide" is a nucleotide sequence such as a nucleic acid fragment. A polynucleotide may be a polymer of RNA or DNA that is single- or double-stranded, that optionally contains synthetic, non-natural or altered nucleotide bases. A polynucleotide in the form of a polymer of DNA may be comprised of one or more segments of cDNA, genomic DNA, synthetic DNA, or mixtures thereof. An isolated polynucleotide of the present invention may include at least one of 60 contiguous nucleotides, preferably at least one of 40 contiguous nucleotides, most preferably one of at least 30 contiguous nucleotides derived from SEQ ID NOs:1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, or the complement of such sequences.

As used herein, "contig" refers to a nucleotide sequence that is assembled from two or more constituent nucleotide sequences that share common or overlapping regions of sequence homology. For example, the nucleotide sequences of two or more nucleic acid fragments can be compared and aligned in order to identify common or overlapping sequences. Where common or overlapping sequences exist between two or more nucleic

acid fragments, the sequences (and thus their corresponding nucleic acid fragments) can be assembled into a single contiguous nucleotide sequence.

As used herein, "substantially similar" refers to nucleic acid fragments wherein changes in one or more nucleotide bases results in substitution of one or more amino acids, but do not affect the functional properties of the polypeptide encoded by the nucleotide sequence. "Substantially similar" also refers to nucleic acid fragments wherein changes in one or more nucleotide bases does not affect the ability of the nucleic acid fragment to mediate alteration of gene expression by gene silencing through for example antisense or co-suppression technology. "Substantially similar" also refers to modifications of the nucleic acid fragments of the instant invention such as deletion or insertion of one or more nucleotides that do not substantially affect the functional properties of the resulting transcript vis-à-vis the ability to mediate gene silencing or alteration of the functional properties of the resulting protein molecule. It is therefore understood that the invention encompasses more than the specific exemplary nucleotide or amino acid sequences and includes functional equivalents thereof.

Substantially similar nucleic acid fragments may be selected by screening nucleic acid fragments representing subfragments or modifications of the nucleic acid fragments of the instant invention, wherein one or more nucleotides are substituted, deleted and/or inserted, for their ability to affect the level of the polypeptide encoded by the unmodified nucleic acid fragment in a plant or plant cell. For example, a substantially similar nucleic acid fragment representing at least one of 30 contiguous nucleotides derived from the instant nucleic acid fragment can be constructed and introduced into a plant or plant cell. The level of the polypeptide encoded by the unmodified nucleic acid fragment present in a plant or plant cell exposed to the substantially similar nucleic fragment can then be compared to the level of the polypeptide in a plant or plant cell that is not exposed to the substantially similar nucleic acid fragment.

For example, it is well known in the art that antisense suppression and co-suppression of gene expression may be accomplished using nucleic acid fragments representing less than the entire coding region of a gene, and by nucleic acid fragments that do not share 100% sequence identity with the gene to be suppressed. Moreover, alterations in a nucleic acid fragment which result in the production of a chemically equivalent amino acid at a given site, but do not effect the functional properties of the encoded polypeptide, are well known in the art. Thus, a codon for the amino acid alanine, a hydrophobic amino acid, may be substituted by a codon encoding another less hydrophobic residue, such as glycine, or a more hydrophobic residue, such as valine, leucine, or isoleucine. Similarly, changes which result in substitution of one negatively charged residue for another, such as aspartic acid for glutamic acid, or one positively charged residue for another, such as lysine for arginine, can also be expected to produce a functionally equivalent product. Nucleotide changes which

result in alteration of the N-terminal and C-terminal portions of the polypeptide molecule would also not be expected to alter the activity of the polypeptide. Each of the proposed modifications is well within the routine skill in the art, as is determination of retention of biological activity of the encoded products. Consequently, an isolated polynucleotide comprising a nucleotide sequence of at least one of 60 (preferably at least one of 40, most preferably at least one of 30) contiguous nucleotides derived from a nucleotide sequence selected from the group consisting of SEQ ID NOs:1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37 and the complement of such nucleotide sequences may be used in methods of selecting an isolated polynucleotide that affects the expression of a polypeptide (vitamin E biosynthetic enzyme) in a host cell. A method of selecting an isolated polynucleotide that affects the level of expression of a polypeptide in a host cell (eukaryotic, such as plant or yeast, prokaryotic such as bacterial, or viral) may comprise the steps of: constructing an isolated polynucleotide of the present invention or an isolated chimeric gene of the present invention; introducing the isolated polynucleotide or the isolated chimeric gene into a host cell; measuring the level a polypeptide in the host cell containing the isolated polynucleotide; and comparing the level of a polypeptide in the host cell containing the isolated polynucleotide with the level of a polypeptide in a host cell that does not contain the isolated polynucleotide.

Moreover, substantially similar nucleic acid fragments may also be characterized by their ability to hybridize. Estimates of such homology are provided by either DNA-DNA or DNA-RNA hybridization under conditions of stringency as is well understood by those skilled in the art (Hames and Higgins, Eds. (1985) *Nucleic Acid Hybridisation*, IRL Press, Oxford, U.K.). Stringency conditions can be adjusted to screen for moderately similar fragments, such as homologous sequences from distantly related organisms, to highly similar fragments, such as genes that duplicate functional enzymes from closely related organisms. Post-hybridization washes determine stringency conditions. One set of preferred conditions uses a series of washes starting with 6X SSC, 0.5% SDS at room temperature for 15 min, then repeated with 2X SSC, 0.5% SDS at 45°C for 30 min, and then repeated twice with 0.2X SSC, 0.5% SDS at 50°C for 30 min. A more preferred set of stringent conditions uses higher temperatures in which the washes are identical to those above except for the temperature of the final two 30 min washes in 0.2X SSC, 0.5% SDS was increased to 60°C. Another preferred set of highly stringent conditions uses two final washes in 0.1X SSC, 0.1% SDS at 65°C.

Substantially similar nucleic acid fragments of the instant invention may also be characterized by the percent identity of the amino acid sequences that they encode to the amino acid sequences disclosed herein, as determined by algorithms commonly employed by those skilled in this art. Suitable nucleic acid fragments (isolated polynucleotides of the present invention) encode polypeptides that are at least about 70% identical, preferably at

least about 80% identical to the amino acid sequences reported herein. Preferred nucleic acid fragments encode amino acid sequences that are at least about 85% identical to the amino acid sequences reported herein. More preferred nucleic acid fragments encode amino acid sequences that are at least about 90% identical to the amino acid sequences reported herein. Most preferred are nucleic acid fragments that encode amino acid sequences that are at least about 95% identical to the amino acid sequences reported herein. Suitable nucleic acid fragments not only have the above homologies but typically encode a polypeptide having at least about 50 amino acids, preferably at least about 100 amino acids, more preferably at least about 150 amino acids, still more preferably at least about 200 amino acids, and most preferably at least about 250 amino acids. Sequence alignments and percent identity calculations were performed using the Megalign program of the LASERGENE bioinformatics computing suite (DNASTAR Inc., Madison, WI). Multiple alignment of the sequences was performed using the Clustal method of alignment (Higgins and Sharp (1989) *CABIOS*. 5:151-153) with the default parameters (GAP PENALTY=10, GAP LENGTH PENALTY=10). Default parameters for pairwise alignments using the Clustal method were KTUPLE 1, GAP PENALTY=3, WINDOW=5 and DIAGONALS SAVED=5.

A "substantial portion" of an amino acid or nucleotide sequence comprises an amino acid or a nucleotide sequence that is sufficient to afford putative identification of the protein or gene that the amino acid or nucleotide sequence comprises. Amino acid and nucleotide sequences can be evaluated either manually by one skilled in the art, or by using computer-based sequence comparison and identification tools that employ algorithms such as BLAST (Basic Local Alignment Search Tool; Altschul et al. (1993) *J. Mol. Biol.* 215:403-410; see also www.ncbi.nlm.nih.gov/BLAST/). In general, a sequence of ten or more contiguous amino acids or thirty or more contiguous nucleotides is necessary in order to putatively identify a polypeptide or nucleic acid sequence as homologous to a known protein or gene. Moreover, with respect to nucleotide sequences, gene-specific oligonucleotide probes comprising 30 or more contiguous nucleotides may be used in sequence-dependent methods of gene identification (e.g., Southern hybridization) and isolation (e.g., *in situ* hybridization of bacterial colonies or bacteriophage plaques). In addition, short oligonucleotides of 12 or more nucleotides may be used as amplification primers in PCR in order to obtain a particular nucleic acid fragment comprising the primers. Accordingly, a "substantial portion" of a nucleotide sequence comprises a nucleotide sequence that will afford specific identification and/or isolation of a nucleic acid fragment comprising the sequence. The instant specification teaches amino acid and nucleotide sequences encoding polypeptides that comprise one or more particular plant proteins. The skilled artisan, having the benefit of the sequences as reported herein, may now use all or a substantial portion of the disclosed sequences for purposes known to those skilled in this art. Accordingly, the instant invention

comprises the complete sequences as reported in the accompanying Sequence Listing, as well as substantial portions of those sequences as defined above.

“Codon degeneracy” refers to divergence in the genetic code permitting variation of the nucleotide sequence without effecting the amino acid sequence of an encoded polypeptide. Accordingly, the instant invention relates to any nucleic acid fragment comprising a nucleotide sequence that encodes all or a substantial portion of the amino acid sequences set forth herein. The skilled artisan is well aware of the “codon-bias” exhibited by a specific host cell in usage of nucleotide codons to specify a given amino acid. Therefore, when synthesizing a nucleic acid fragment for improved expression in a host cell, it is desirable to design the nucleic acid fragment such that its frequency of codon usage approaches the frequency of preferred codon usage of the host cell.

“Synthetic nucleic acid fragments” can be assembled from oligonucleotide building blocks that are chemically synthesized using procedures known to those skilled in the art. These building blocks are ligated and annealed to form larger nucleic acid fragments which may then be enzymatically assembled to construct the entire desired nucleic acid fragment. “Chemically synthesized”, as related to nucleic acid fragment, means that the component nucleotides were assembled *in vitro*. Manual chemical synthesis of nucleic acid fragments may be accomplished using well established procedures, or automated chemical synthesis can be performed using one of a number of commercially available machines. Accordingly, the nucleic acid fragments can be tailored for optimal gene expression based on optimization of nucleotide sequence to reflect the codon bias of the host cell. The skilled artisan appreciates the likelihood of successful gene expression if codon usage is biased towards those codons favored by the host. Determination of preferred codons can be based on a survey of genes derived from the host cell where sequence information is available.

“Gene” refers to a nucleic acid fragment that expresses a specific protein, including regulatory sequences preceding (5' non-coding sequences) and following (3' non-coding sequences) the coding sequence. “Native gene” refers to a gene as found in nature with its own regulatory sequences. “Chimeric gene” refers any gene that is not a native gene, comprising regulatory and coding sequences that are not found together in nature. Accordingly, a chimeric gene may comprise regulatory sequences and coding sequences that are derived from different sources, or regulatory sequences and coding sequences derived from the same source, but arranged in a manner different than that found in nature. “Endogenous gene” refers to a native gene in its natural location in the genome of an organism. A “foreign” gene refers to a gene not normally found in the host organism, but that is introduced into the host organism by gene transfer. Foreign genes can comprise native genes inserted into a non-native organism, or chimeric genes. A “transgene” is a gene that has been introduced into the genome by a transformation procedure.

“Coding sequence” refers to a nucleotide sequence that codes for a specific amino acid sequence. “Regulatory sequences” refer to nucleotide sequences located upstream (5' non-coding sequences), within, or downstream (3' non-coding sequences) of a coding sequence, and which influence the transcription, RNA processing or stability, or translation of the associated coding sequence. Regulatory sequences may include promoters, translation leader sequences, introns, and polyadenylation recognition sequences.

“Promoter” refers to a nucleotide sequence capable of controlling the expression of a coding sequence or functional RNA. In general, a coding sequence is located 3' to a promoter sequence. The promoter sequence consists of proximal and more distal upstream elements, the latter elements often referred to as enhancers. Accordingly, an “enhancer” is a nucleotide sequence which can stimulate promoter activity and may be an innate element of the promoter or a heterologous element inserted to enhance the level or tissue-specificity of a promoter. Promoters may be derived in their entirety from a native gene, or be composed of different elements derived from different promoters found in nature, or even comprise synthetic nucleotide segments. It is understood by those skilled in the art that different promoters may direct the expression of a gene in different tissues or cell types, or at different stages of development, or in response to different environmental conditions. Promoters which cause a nucleic acid fragment to be expressed in most cell types at most times are commonly referred to as “constitutive promoters”. New promoters of various types useful in plant cells are constantly being discovered; numerous examples may be found in the compilation by Okamuro and Goldberg (1989) *Biochemistry of Plants* 15:1-82. It is further recognized that since in most cases the exact boundaries of regulatory sequences have not been completely defined, nucleic acid fragments of different lengths may have identical promoter activity.

The “translation leader sequence” refers to a nucleotide sequence located between the promoter sequence of a gene and the coding sequence. The translation leader sequence is present in the fully processed mRNA upstream of the translation start sequence. The translation leader sequence may affect processing of the primary transcript to mRNA, mRNA stability or translation efficiency. Examples of translation leader sequences have been described (Turner and Foster (1995) *Mol. Biotechnol.* 3:225-236).

The “3' non-coding sequences” refer to nucleotide sequences located downstream of a coding sequence and include polyadenylation recognition sequences and other sequences encoding regulatory signals capable of affecting mRNA processing or gene expression. The polyadenylation signal is usually characterized by affecting the addition of polyadenylic acid tracts to the 3' end of the mRNA precursor. The use of different 3' non-coding sequences is exemplified by Ingelbrecht et al. (1989) *Plant Cell* 1:671-680.

“RNA transcript” refers to the product resulting from RNA polymerase-catalyzed transcription of a DNA sequence. When the RNA transcript is a perfect complementary

copy of the DNA sequence, it is referred to as the primary transcript or it may be a RNA sequence derived from posttranscriptional processing of the primary transcript and is referred to as the mature RNA. "Messenger RNA (mRNA)" refers to the RNA that is without introns and that can be translated into polypeptide by the cell. "cDNA" refers to a double-stranded DNA that is complementary to and derived from mRNA. "Sense" RNA refers to an RNA transcript that includes the mRNA and so can be translated into a polypeptide by the cell. "Antisense RNA" refers to an RNA transcript that is complementary to all or part of a target primary transcript or mRNA and that blocks the expression of a target gene (see U.S. Patent No. 5,107,065, incorporated herein by reference). The complementarity of an antisense RNA may be with any part of the specific nucleotide sequence, i.e., at the 5' non-coding sequence, 3' non-coding sequence, introns, or the coding sequence. "Functional RNA" refers to sense RNA, antisense RNA, ribozyme RNA, or other RNA that may not be translated but yet has an effect on cellular processes.

The term "operably linked" refers to the association of two or more nucleic acid fragments on a single nucleic acid fragment so that the function of one is affected by the other. For example, a promoter is operably linked with a coding sequence when it is capable of affecting the expression of that coding sequence (i.e., that the coding sequence is under the transcriptional control of the promoter). Coding sequences can be operably linked to regulatory sequences in sense or antisense orientation.

The term "expression", as used herein, refers to the transcription and stable accumulation of sense (mRNA) or antisense RNA derived from the nucleic acid fragment of the invention. Expression may also refer to translation of mRNA into a polypeptide. "Antisense inhibition" refers to the production of antisense RNA transcripts capable of suppressing the expression of the target protein. "Overexpression" refers to the production of a gene product in transgenic organisms that exceeds levels of production in normal or non-transformed organisms. "Co-suppression" refers to the production of sense RNA transcripts capable of suppressing the expression of identical or substantially similar foreign or endogenous genes (U.S. Patent No. 5,231,020, incorporated herein by reference).

"Altered levels" refers to the production of gene product(s) in transgenic organisms in amounts or proportions that differ from that of normal or non-transformed organisms.

"Mature" protein refers to a post-translationally processed polypeptide; i.e., one from which any pre- or propeptides present in the primary translation product have been removed. "Precursor" protein refers to the primary product of translation of mRNA; i.e., with pre- and propeptides still present. Pre- and propeptides may be but are not limited to intracellular localization signals.

A "chloroplast transit peptide" is an amino acid sequence which is translated in conjunction with a protein and directs the protein to the chloroplast or other plastid types present in the cell in which the protein is made. "Chloroplast transit sequence" refers to a

nucleotide sequence that encodes a chloroplast transit peptide. A "signal peptide" is an amino acid sequence which is translated in conjunction with a protein and directs the protein to the secretory system (Chrispeels (1991) *Ann. Rev. Plant Phys. Plant Mol. Biol.* 42:21-53). If the protein is to be directed to a vacuole, a vacuolar targeting signal (*supra*) can further be added, or if to the endoplasmic reticulum, an endoplasmic reticulum retention signal (*supra*) may be added. If the protein is to be directed to the nucleus, any signal peptide present should be removed and instead a nuclear localization signal included (Raikhel (1992) *Plant Phys.* 100:1627-1632).

"Transformation" refers to the transfer of a nucleic acid fragment into the genome of a host organism, resulting in genetically stable inheritance. Host organisms containing the transformed nucleic acid fragments are referred to as "transgenic" organisms. Examples of methods of plant transformation include *Agrobacterium*-mediated transformation (De Blaere et al. (1987) *Meth. Enzymol.* 143:277) and particle-accelerated or "gene gun" transformation technology (Klein et al. (1987) *Nature (London)* 327:70-73; U.S. Patent No. 4,945,050, incorporated herein by reference).

Standard recombinant DNA and molecular cloning techniques used herein are well known in the art and are described more fully in Sambrook et al. *Molecular Cloning: A Laboratory Manual*; Cold Spring Harbor Laboratory Press: Cold Spring Harbor, 1989 (hereinafter "Maniatis").

Nucleic acid fragments encoding at least a portion of several vitamin E biosynthetic enzymes have been isolated and identified by comparison of random plant cDNA sequences to public databases containing nucleotide and protein sequences using the BLAST algorithms well known to those skilled in the art. The nucleic acid fragments of the instant invention may be used to isolate cDNAs and genes encoding homologous proteins from the same or other plant species. Isolation of homologous genes using sequence-dependent protocols is well known in the art. Examples of sequence-dependent protocols include, but are not limited to, methods of nucleic acid hybridization, and methods of DNA and RNA amplification as exemplified by various uses of nucleic acid amplification technologies (e.g., polymerase chain reaction, ligase chain reaction).

For example, genes encoding other 4-hydroxyphenylpyruvate dioxygenases or gamma tocopherol methyltransferases, either as cDNAs or genomic DNAs, could be isolated directly by using all or a portion of the instant nucleic acid fragments as DNA hybridization probes to screen libraries from any desired plant employing methodology well known to those skilled in the art. Specific oligonucleotide probes based upon the instant nucleic acid sequences can be designed and synthesized by methods known in the art (Maniatis).

Moreover, the entire sequences can be used directly to synthesize DNA probes by methods known to the skilled artisan such as random primer DNA labeling, nick translation, or end-labeling techniques, or RNA probes using available *in vitro* transcription systems. In

addition, specific primers can be designed and used to amplify a part or all of the instant sequences. The resulting amplification products can be labeled directly during amplification reactions or labeled after amplification reactions, and used as probes to isolate full length cDNA or genomic fragments under conditions of appropriate stringency.

- 5 In addition, two short segments of the instant nucleic acid fragments may be used in polymerase chain reaction protocols to amplify longer nucleic acid fragments encoding homologous genes from DNA or RNA. The polymerase chain reaction may also be performed on a library of cloned nucleic acid fragments wherein the sequence of one primer is derived from the instant nucleic acid fragments, and the sequence of the other primer takes
- 10 advantage of the presence of the polyadenylic acid tracts to the 3' end of the mRNA precursor encoding plant genes. Alternatively, the second primer sequence may be based upon sequences derived from the cloning vector. For example, the skilled artisan can follow the RACE protocol (Frohman et al. (1988) *Proc. Natl. Acad. Sci. USA* 85:8998-9002) to generate cDNAs by using PCR to amplify copies of the region between a single point in the
- 15 transcript and the 3' or 5' end. Primers oriented in the 3' and 5' directions can be designed from the instant sequences. Using commercially available 3' RACE or 5' RACE systems (BRL), specific 3' or 5' cDNA fragments can be isolated (Ohara et al. (1989) *Proc. Natl. Acad. Sci. USA* 86:5673-5677; Loh et al. (1989) *Science* 243:217-220). Products generated by the 3' and 5' RACE procedures can be combined to generate full-length cDNAs (Frohman
- 20 and Martin (1989) *Techniques* 1:165). Consequently, a polynucleotide comprising a nucleotide sequence of at least one of 60 (preferably one of at least 40, most preferably one of at least 30) contiguous nucleotides derived from a nucleotide sequence selected from the group consisting of SEQ ID NOs:1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37 and the complement of such nucleotide sequences may be used in such methods to
- 25 obtain a nucleic acid fragment encoding a substantial portion of an amino acid sequence of a polypeptide. The present invention relates to a method of obtaining a nucleic acid fragment encoding a substantial portion of a polypeptide of a gene (such as gamma tocopherol methyltransferase or 4-phenylpyruvate dioxygenase) preferably a substantial portion of a plant polypeptide of a gene, comprising the steps of: synthesizing an oligonucleotide primer
- 30 comprising a nucleotide sequence of at least one of 60 (preferably at least one of 40, most preferably at least one of 30) contiguous nucleotides derived from a nucleotide sequence selected from the group consisting of SEQ ID NOs:1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, and the complement of such nucleotide sequences; and amplifying a nucleic acid fragment (preferably a cDNA inserted in a cloning vector) using the
- 35 oligonucleotide primer. The amplified nucleic acid fragment preferably will encode a portion of a polypeptide (gamma tocopherol methyltransferase or 4-phenylpyruvate dioxygenase).

Availability of the instant nucleotide and deduced amino acid sequences facilitates immunological screening of cDNA expression libraries. Synthetic peptides representing portions of the instant amino acid sequences may be synthesized. These peptides can be used to immunize animals to produce polyclonal or monoclonal antibodies with specificity
5 for peptides or proteins comprising the amino acid sequences. These antibodies can be then be used to screen cDNA expression libraries to isolate full-length cDNA clones of interest (Lerner (1984) *Adv. Immunol.* 36:1-34; Maniatis).

The nucleic acid fragments of the instant invention may be used to create transgenic plants in which the disclosed polypeptides are present at higher or lower levels than normal
10 or in cell types or developmental stages in which they are not normally found. This would have the effect of altering the level of homogénistate in those cells. Overexpression of 4-hydroxyphenylpyruvate dioxygenase should result in a larger accumulation of homogénistate which may be used by gamma tocopherol methyltransferase to produce vitamin E. Since mammals can not synthesize tocopherols, the enzymes described herein
15 may be used for the discovery of new herbicides.

Overexpression of the proteins of the instant invention may be accomplished by first constructing a chimeric gene in which the coding region is operably linked to a promoter capable of directing expression of a gene in the desired tissues at the desired stage of development. For reasons of convenience, the chimeric gene may comprise promoter
20 sequences and translation leader sequences derived from the same genes. 3' Non-coding sequences encoding transcription termination signals may also be provided. The instant chimeric gene may also comprise one or more introns in order to facilitate gene expression.

Plasmid vectors comprising the instant chimeric gene can then be constructed. The choice of plasmid vector is dependent upon the method that will be used to transform host
25 plants. The skilled artisan is well aware of the genetic elements that must be present on the plasmid vector in order to successfully transform, select and propagate host cells containing the chimeric gene. The skilled artisan will also recognize that different independent transformation events will result in different levels and patterns of expression (Jones et al. (1985) *EMBO J.* 4:2411-2418; De Almeida et al. (1989) *Mol. Gen. Genetics* 218:78-86), and
30 thus that multiple events must be screened in order to obtain lines displaying the desired expression level and pattern. Such screening may be accomplished by Southern analysis of DNA, Northern analysis of mRNA expression, Western analysis of protein expression, or phenotypic analysis.

For some applications it may be useful to direct the instant polypeptides to different
35 cellular compartments, or to facilitate its secretion from the cell. It is thus envisioned that the chimeric gene described above may be further supplemented by directing the coding sequence to encode the instant polypeptides with appropriate intracellular targeting sequences such as transit sequences (Keegstra (1989) *Cell* 56:247-253), signal sequences or

sequences encoding endoplasmic reticulum localization (Chrispeels (1991) *Ann. Rev. Plant Phys. Plant Mol. Biol.* 42:21-53), or nuclear localization signals (Raikhel (1992) *Plant Phys.* 100:1627-1632) with or without removing targeting sequences that are already present. While the references cited give examples of each of these, the list is not exhaustive and more
5 targeting signals of use may be discovered in the future.

It may also be desirable to reduce or eliminate expression of genes encoding the instant polypeptides in plants for some applications. In order to accomplish this, a chimeric gene designed for co-suppression of the instant polypeptide can be constructed by linking a gene or gene fragment encoding that polypeptide to plant promoter sequences.

10 Alternatively, a chimeric gene designed to express antisense RNA for all or part of the instant nucleic acid fragment can be constructed by linking the gene or gene fragment in reverse orientation to plant promoter sequences. Either the co-suppression or antisense chimeric genes could be introduced into plants via transformation wherein expression of the corresponding endogenous genes are reduced or eliminated.

15 Molecular genetic solutions to the generation of plants with altered gene expression have a decided advantage over more traditional plant breeding approaches. Changes in plant phenotypes can be produced by specifically inhibiting expression of one or more genes by antisense inhibition or cosuppression (U.S. Patent Nos. 5,190,931, 5,107,065 and 5,283,323). An antisense or cosuppression construct would act as a dominant negative
20 regulator of gene activity. While conventional mutations can yield negative regulation of gene activity these effects are most likely recessive. The dominant negative regulation available with a transgenic approach may be advantageous from a breeding perspective. In addition, the ability to restrict the expression of specific phenotype to the reproductive tissues of the plant by the use of tissue specific promoters may confer agronomic advantages
25 relative to conventional mutations which may have an effect in all tissues in which a mutant gene is ordinarily expressed.

The person skilled in the art will know that special considerations are associated with the use of antisense or cosuppression technologies in order to reduce expression of particular genes. For example, the proper level of expression of sense or antisense genes may require
30 the use of different chimeric genes utilizing different regulatory elements known to the skilled artisan. Once transgenic plants are obtained by one of the methods described above, it will be necessary to screen individual transgenics for those that most effectively display the desired phenotype. Accordingly, the skilled artisan will develop methods for screening large numbers of transformants. The nature of these screens will generally be chosen on
35 practical grounds, and is not an inherent part of the invention. For example, one can screen by looking for changes in gene expression by using antibodies specific for the protein encoded by the gene being suppressed, or one could establish assays that specifically measure enzyme activity. A preferred method will be one which allows large numbers of

samples to be processed rapidly, since it will be expected that a large number of transformants will be negative for the desired phenotype.

5 The instant polypeptides (or portions thereof) may be produced in heterologous host cells, particularly in the cells of microbial hosts, and can be used to prepare antibodies to these proteins by methods well known to those skilled in the art. The antibodies are useful for detecting the polypeptides of the instant invention *in situ* in cells or *in vitro* in cell extracts. Preferred heterologous host cells for production of the instant polypeptides are microbial hosts. Microbial expression systems and expression vectors containing regulatory sequences that direct high level expression of foreign proteins are well known to those skilled in the art. Any of these could be used to construct a chimeric gene for production of the instant polypeptides. This chimeric gene could then be introduced into appropriate microorganisms via transformation to provide high level expression of the encoded vitamin E biosynthetic enzyme. An example of a vector for high level expression of the instant polypeptides in a bacterial host is provided (Example 7).

15 Additionally, the instant polypeptides can be used as a targets to facilitate design and/or identification of inhibitors of those enzymes that may be useful as herbicides. This is desirable because the polypeptides described herein catalyze various steps in vitamin E biosynthesis. Accordingly, inhibition of the activity of one or more of the enzymes described herein could lead to inhibition of plant growth. Thus, the instant polypeptides could be appropriate for new herbicide discovery and design.

20 All or a substantial portion of the nucleic acid fragments of the instant invention may also be used as probes for genetically and physically mapping the genes that they are a part of, and as markers for traits linked to those genes. Such information may be useful in plant breeding in order to develop lines with desired phenotypes. For example, the instant nucleic acid fragments may be used as restriction fragment length polymorphism (RFLP) markers. Southern blots (Maniatis) of restriction-digested plant genomic DNA may be probed with the nucleic acid fragments of the instant invention. The resulting banding patterns may then be subjected to genetic analyses using computer programs such as MapMaker (Lander et al. (1987) *Genomics* 1:174-181) in order to construct a genetic map. In addition, the nucleic acid fragments of the instant invention may be used to probe Southern blots containing restriction endonuclease-treated genomic DNAs of a set of individuals representing parent and progeny of a defined genetic cross. Segregation of the DNA polymorphisms is noted and used to calculate the position of the instant nucleic acid sequence in the genetic map previously obtained using this population (Botstein et al. (1980) *Am. J. Hum. Genet.* 32:314-331).

35 The production and use of plant gene-derived probes for use in genetic mapping is described in Bernatzky and Tanksley (1986) *Plant Mol. Biol. Reporter* 4:37-41. Numerous publications describe genetic mapping of specific cDNA clones using the methodology

outlined above or variations thereof. For example, F2 intercross populations, backcross populations, randomly mated populations, near isogenic lines, and other sets of individuals may be used for mapping. Such methodologies are well known to those skilled in the art.

5 Nucleic acid probes derived from the instant nucleic acid sequences may also be used for physical mapping (i.e., placement of sequences on physical maps; see Hoheisel et al. In: *Nonmammalian Genomic Analysis: A Practical Guide*, Academic press 1996, pp. 319-346, and references cited therein).

10 In another embodiment, nucleic acid probes derived from the instant nucleic acid sequences may be used in direct fluorescence *in situ* hybridization (FISH) mapping (Trask (1991) *Trends Genet.* 7:149-154). Although current methods of FISH mapping favor use of large clones (several to several hundred KB; see Laan et al. (1995) *Genome Res.* 5:13-20), improvements in sensitivity may allow performance of FISH mapping using shorter probes.

A variety of nucleic acid amplification-based methods of genetic and physical mapping may be carried out using the instant nucleic acid sequences. Examples include
15 allele-specific amplification (Kazazian (1989) *J. Lab. Clin. Med.* 11:95-96), polymorphism of PCR-amplified fragments (CAPS; Sheffield et al. (1993) *Genomics* 16:325-332), allele-specific ligation (Landegren et al. (1988) *Science* 241:1077-1080), nucleotide extension reactions (Sokolov (1990) *Nucleic Acid Res.* 18:3671), Radiation Hybrid Mapping (Walter et al. (1997) *Nat. Genet.* 7:22-28) and Happy Mapping (Dear and Cook (1989) *Nucleic Acid*
20 *Res.* 17:6795-6807). For these methods, the sequence of a nucleic acid fragment is used to design and produce primer pairs for use in the amplification reaction or in primer extension reactions. The design of such primers is well known to those skilled in the art. In methods employing PCR-based genetic mapping, it may be necessary to identify DNA sequence differences between the parents of the mapping cross in the region corresponding to the
25 instant nucleic acid sequence. This, however, is generally not necessary for mapping methods.

Loss of function mutant phenotypes may be identified for the instant cDNA clones either by targeted gene disruption protocols or by identifying specific mutants for these genes contained in a maize population carrying mutations in all possible genes (Ballinger
30 and Benzer (1989) *Proc. Natl. Acad. Sci USA* 86:9402-9406; Koes et al. (1995) *Proc. Natl. Acad. Sci USA* 92:8149-8153; Bensen et al. (1995) *Plant Cell* 7:75-84). The latter approach may be accomplished in two ways. First, short segments of the instant nucleic acid fragments may be used in polymerase chain reaction protocols in conjunction with a mutation tag sequence primer on DNAs prepared from a population of plants in which
35 Mutator transposons or some other mutation-causing DNA element has been introduced (see Bensen, *supra*). The amplification of a specific DNA fragment with these primers indicates the insertion of the mutation tag element in or near the plant gene encoding the instant polypeptides. Alternatively, the instant nucleic acid fragment may be used as a

hybridization probe against PCR amplification products generated from the mutation population using the mutation tag sequence primer in conjunction with an arbitrary genomic site primer, such as that for a restriction enzyme site-anchored synthetic adaptor. With either method, a plant containing a mutation in the endogenous gene encoding the instant polypeptides can be identified and obtained. This mutant plant can then be used to determine or confirm the natural function of the instant polypeptides disclosed herein.

EXAMPLES

The present invention is further defined in the following Examples, in which all parts and percentages are by weight and degrees are Celsius, unless otherwise stated. It should be understood that these Examples, while indicating preferred embodiments of the invention, are given by way of illustration only. From the above discussion and these Examples, one skilled in the art can ascertain the essential characteristics of this invention, and without departing from the spirit and scope thereof, can make various changes and modifications of the invention to adapt it to various usages and conditions.

EXAMPLE 1

Composition of cDNA Libraries; Isolation and Sequencing of cDNA Clones

cDNA libraries representing mRNAs from various corn, rice, soybean, *Vernonia*, and wheat tissues were prepared. The characteristics of the libraries are described below.

TABLE 2

cDNA Libraries from Corn, Rice, Soybean, *Vernonia*, and Wheat

Library	Tissue	Clone
cr1n	Corn Root From 7 Day Old Seedlings*	cr1n.pk0155.d1
cr1n	Corn Root From 7 Day Old Seedlings*	cr1n.pk0179.f10
cs1	Corn Leaf Sheath From 5 Week Old Plant	cs1.pk0065.e1
cs1	Corn Leaf Sheath From 5 Week Old Plant	cs1.pk0065.f2
cta1n	Corn Tassel*	cta1n.pk0031.d2
ncs	<i>Catalpa speciosa</i> Developing Seed	ncs.pk0012.g1
p0023	Corn Leaf From Plant Transformed with Gene M1C07 (leucine-rich repeat) Which Induces Resistance Prior to Genetic Lesion Formation. Harvested About One Month After Planting in Green House*	p0060.coran49r
p0103	Corn Tassel Shoots(0.1-1.4 cm)*	p0103.ciaay86r
rl0n	Rice 15 Day Old Leaf*	rl0n.pk085.e11
rl0n	Rice 15 Day Old Leaf*	rl0n.pk099.d24
rlr12	Rice Leaf 15 Days After Germination, 12 Hours After Infection of Strain <i>Magaporthe grisea</i> 4360-R-62 (AVR2-YAMO); Resistant	rlr12.pk0019.g5

Library	Tissue	Clone
rlr12	Rice Leaf 15 Days After Germination, 12 Hours After Infection of Strain <i>Magaporthe grisea</i> 4360-R-62 (AVR2-YAMO); Resistant	rlr12.pk0025.e4
rls72	Rice Leaf 15 Days After Germination, 72 Hours After Infection of Strain <i>Magaporthe grisea</i> 4360-R-67 (AVR2-YAMO); Susceptible	rls72.pk0010.g3
sah1c	Soybean Sprayed With Authority Herbicide	sah1c.pk001.k8
sah1c	Soybean Sprayed With Authority Herbicide	sah1c.pk004.g2
sfl1	Soybean Immature Flower	sfl1.pk126.n15
sgc4c	Soybean Cotyledon 14-21 Days After Germination (1/4 yellow)	sgc4c.pk001.p2
sgc5c	Soybean (Cotyledon 15-24 Days After Germination (3/4 yellow)	sgc5c.pk001.j9
sgc5c	Soybean (Cotyledon 15-24 Days After Germination (3/4 yellow)	sgc5c.pk001.m23
sgs1c	Soybean Seeds 4 Hours After Germination	sgs1c.pk002.a8
sgs1c	Soybean Seeds 4 Hours After Germination	sgs1c.pk003.o5
vs1	<i>Vernonia</i> Seed Stage 1	vs1.pk0015.b2
wdk4c	Wheat Developing Kernel, 21 Days After Anthesis	wdk4c.pk006.m9
wle1n	Wheat Leaf From 7 Day Old Etiolated Seedling*	wle1n.pk0065.h9
wlm0	Wheat Seedlings 0 Hour After Inoculation With <i>Erysiphe graminis f. sp tritici</i>	wlm0.pk0035.d5
wlmk1	Wheat Seedlings 1 Hour After Inoculation With <i>Erysiphe graminis f. sp tritici</i> and Treatment With Herbicide**	wlmk1.pk0013.g11
wlmk1	Wheat Seedlings 1 Hour After Inoculation With <i>Erysiphe graminis f. sp tritici</i> and Treatment With Herbicide**	wlmk1.pk0019.e2
wlmk1	Wheat Seedlings 1 Hour After Inoculation With <i>Erysiphe graminis f. sp tritici</i> and Treatment With Herbicide**	wlmk1.pk0021.h10
wr1	Wheat Root From 7 Day Old Seedling	wr1.pk0077.fl
wr1	Wheat Root From 7 Day Old Seedling	wr1.pk177.b11
wre1n	Wheat Root From 7 Day Old Etiolated Seedling*	wre1n.pk0111.d6

* These libraries were normalized essentially as described in U.S. Patent No. 5,482,845, incorporated herein by reference.

5 ** Application of 6-iodo-2-propoxy-3-propyl-4(3H)-quinazolinone; synthesis and methods of using this compound are described in USSN 08/545,827, incorporated herein by reference.

10 cDNA libraries may be prepared by any one of many methods available. For example, the cDNAs may be introduced into plasmid vectors by first preparing the cDNA libraries in Uni-ZAP™ XR vectors according to the manufacturer's protocol (Stratagene

Cloning Systems, La Jolla, CA). The Uni-ZAP™ XR libraries are converted into plasmid libraries according to the protocol provided by Stratagene. Upon conversion, cDNA inserts will be contained in the plasmid vector pBluescript. In addition, the cDNAs may be introduced directly into precut Bluescript II SK(+) vectors (Stratagene) using T4 DNA ligase (New England Biolabs), followed by transfection into DH10B cells according to the manufacturer's protocol (GIBCO BRL Products). Once the cDNA inserts are in plasmid vectors, plasmid DNAs are prepared from randomly picked bacterial colonies containing recombinant pBluescript plasmids, or the insert cDNA sequences are amplified via polymerase chain reaction using primers specific for vector sequences flanking the inserted cDNA sequences. Amplified insert DNAs or plasmid DNAs are sequenced in dye-primer sequencing reactions to generate partial cDNA sequences (expressed sequence tags or "ESTs"; see Adams et al., (1991) *Science* 252:1651-1656). The resulting ESTs are analyzed using a Perkin Elmer Model 377 fluorescent sequencer.

EXAMPLE 2

Identification of cDNA Clones

cDNA clones encoding vitamin E biosynthetic enzymes were identified by conducting BLAST (Basic Local Alignment Search Tool; Altschul et al. (1993) *J. Mol. Biol.* 215:403-410; see also www.ncbi.nlm.nih.gov/BLAST/) searches for similarity to sequences contained in the BLAST "nr" database (comprising all non-redundant GenBank CDS translations, sequences derived from the 3-dimensional structure Brookhaven Protein Data Bank, the last major release of the SWISS-PROT protein sequence database, EMBL, and DDBJ databases). The cDNA sequences obtained in Example 1 were analyzed for similarity to all publicly available DNA sequences contained in the "nr" database using the BLASTN algorithm provided by the National Center for Biotechnology Information (NCBI). The DNA sequences were translated in all reading frames and compared for similarity to all publicly available protein sequences contained in the "nr" database using the BLASTX algorithm (Gish and States (1993) *Nat. Genet.* 3:266-272) provided by the NCBI. For convenience, the P-value (probability) of observing a match of a cDNA sequence to a sequence contained in the searched databases merely by chance as calculated by BLAST are reported herein as "pLog" values, which represent the negative of the logarithm of the reported P-value. Accordingly, the greater the pLog value, the greater the likelihood that the cDNA sequence and the BLAST "hit" represent homologous proteins.

EXAMPLE 3

Characterization of cDNA Clones Encoding

Gamma Tocopherol Methyltransferase

The BLASTX search using the EST sequences from clones listed in Table 3 revealed similarity of the polypeptides encoded by the cDNAs to gamma tocopherol methyltransferase from *Synechocystis* sp. (NCBI General Identifier No. 1001725). Shown in

Table 3 are the BLAST results for individual ESTs ("EST"), the sequences of the entire cDNA inserts comprising the indicated cDNA clones ("FIS"), contigs assembled from two or more ESTs ("Contig"), contigs assembled from an FIS and one or more ESTs ("Contig*"), or sequences encoding the entire protein derived from an FIS, a contig, or an FIS and PCR ("CGS"):

TABLE 3

BLAST Results for Sequences Encoding Polypeptides Homologous to gamma Tocopherol Methyltransferase

Clone	Status	BLAST pLog Score 1001725
Contig of: crln.pk0179.f10 cs1.pk0065.e1 cs1.pk0065.f2 ctaln.pk0031.d2	Contig	49.70
rl0n.pk085.e11	EST	15.10
Contig of: rl0n.pk099.d24 rls72.pk0010.g3	Contig	8.30
Contig of: sah1c.pk001.k8 sgc5c.pk001.m23 sah1c.pk004.g2:fis	CGS	64.22
Contig of: wr1.pk0077.f1 wr1.pk177.b11 wle1n.pk0065.h9 wre1n.pk0111.d6	Contig	57.00

10

Further sequencing and searching of the DuPont proprietary database allowed for the assembly of longer sequences. The BLASTX search using the nucleotide sequences from clones listed in Table 4 revealed similarity of the polypeptides encoded by the cDNAs to gamma tocopherol methyltransferase from *Arabidopsis thaliana* (NCBI General Identifier No. 4106538). Shown in Table 4 are the BLAST results for individual ESTs ("EST") or sequences encoding the entire protein derived from the sequences of the entire cDNA inserts comprising the indicated cDNA clones ("FIS"), contigs assembled from an FIS and one or more ESTs, or an FIS and PCR ("CGS"):

15

TABLE 4

BLAST Results for Sequences Encoding Polypeptides Homologous
to gamma Tocopherol Methyltransferase

Clone	Status	BLAST pLog Score 4106538
Contig of: crln.pk0179.f10:fis csl.pk0065.f2 ctaln.pk0031.d2 p0060.coran49r p0103.ciaay86r	CGS	118.00
rls72.pk0010.g3-3'	EST*	81.30
rls72.pk0010.g3-5'	EST*	11.00
sahlc.pk004.g2	CGS	125.00
wrl.pk0077.f1:fis	CGS	120.00

5 *These sequences are derived from partial full-insert sequencing of clone rls72.pk0010.g3
and correspond to 5'-end and 3'-end sequences.

Figure 1 presents an alignment of the amino acid sequences set forth in SEQ ID
NOs:8, 22, 24, 26, 28, and 30 and the *Synechocystis sp.* and *Arabidopsis thaliana* sequences
(SEQ ID NO:39 and SEQ ID NO:40, respectively). The data in Table 5 represents a
10 calculation of the percent identity of the amino acid sequences set forth in SEQ ID NOs:8,
22, 24, 26, 28, and 30 and the *Synechocystis sp.* and *Arabidopsis thaliana* sequences (SEQ
ID NO:39 and SEQ ID NO:40, respectively).

TABLE 5

15 Percent Identity of Amino Acid Sequences Deduced From the Nucleotide
Sequences of cDNA Clones Encoding Polypeptides Homologous
to gamma Tocopherol Methyltransferase

SEQ ID NO.	Percent Identity to	
	1001725	4106538
8	37.9	56.9
22	42.6	58.6
24	42.8	67.6
26	23.8	33.3
28	41.3	62.6
30	43.2	59.5

20 Sequence alignments and percent identity calculations were performed using the
Megalign program of the LASERGENE bioinformatics computing suite (DNASTAR Inc.,
Madison, WI). Multiple alignment of the sequences was performed using the Clustal
method of alignment (Higgins and Sharp (1989) *CABIOS*. 5:151-153) with the default

parameters (GAP PENALTY=10, GAP LENGTH PENALTY=10). Default parameters for pairwise alignments using the Clustal method were KTUPLE 1, GAP PENALTY=3, WINDOW=5 and DIAGONALS SAVED=5. Sequence alignments and BLAST scores and probabilities indicate that the nucleic acid fragments comprising the instant cDNA clones encode one entire corn, two entire soybean, one entire wheat, and two fragments corresponding to an almost entire rice gamma tocopherol methyltransferase. These sequences represent the first corn, rice, soybean, and wheat sequences encoding gamma tocopherol methyltransferase.

EXAMPLE 4

Characterization of cDNA Clones Encoding 4-Hydroxyphenylpyruvate Dioxygenase

The BLASTX search using the EST sequences from clones listed in Table 6 revealed similarity of the polypeptides encoded by the cDNAs to 4-hydroxyphenylpyruvate dioxygenase from *Hordeum vulgare* (NCBI General Identifier No. 3334222), *Daucus carota* (NCBI General Identifier No. 3334219) and *Arabidopsis thaliana* (NCBI General Identifier No. 3334223). Shown in Table 6 are the BLAST results for individual ESTs ("EST") or contigs assembled from two or more ESTs ("Contig"):

TABLE 6
BLAST Results for Sequences Encoding Polypeptides Homologous
to 4-Hydroxyphenylpyruvate Dioxygenase

Clone	Status	BLAST pLog Score		
		3334222	3334219	3334223
rlr12.pk0019.g5	EST	26.70	17.70	18.00
rlr12.pk0025.e4	EST	58.52	50.40	49.40
Contig of: sgc5c.pk001.j9 sgs1c.pk002.a8 sfl1.pk126.n15 sgs1c.pk003.o5 sgc4c.pk001.p2	Contig	55.00	81.30	107.00
vs1.pk0015.b2	EST	39.40	51.70	52.05
Contig of: wdk4c.pk006.m9 wlmk1.pk0021.h1 wlmk1.pk0019.e2 wlm0.pk0035.d5 wlmk1.pk0013.g1	Contig	176.00	102.00	97.00

Further sequencing and searching of the DuPont proprietary database allowed the assembly of longer sequences and identification of 4-hydroxyphenylpyruvate dioxygenase in

other species. The BLASTX search using the nucleotide sequences from clones listed in Table 7 revealed similarity of the polypeptides encoded by the cDNAs to 4-hydroxyphenylpyruvate dioxygenase from *Hordeum vulgare* (NCBI General Identifier No. 3334222), *Daucus carota* (NCBI General Identifier No. 3334219) and *Arabidopsis thaliana* (NCBI General Identifier No. 3334223). Shown in Table 7 are the BLAST results for the sequences of the entire cDNA inserts comprising the indicated cDNA clones ("FIS") or sequences encoding the entire protein derived from an FIS, or an FIS and PCR ("CGS"):

TABLE 7

BLAST Results for Sequences Encoding Polypeptides Homologous to 4-Hydroxyphenylpyruvate Dioxygenase

Clone	Status	BLAST pLog Score		
		3334222	3334219	3334223
ncs.pk0012.g1:fis	CGS	147.00	254.00	177.00
rlr12.pk0025.e4:fis	FIS	117.00	102.00	100.00
sgc5c.pk001.j9:fis	CGS	149.00	>254.00	>254.00
wdk4c.pk006.m9:fis	CGS	>254.00	155.00	151.00

Figure 2 presents an alignment of the amino acid sequences set forth in SEQ ID NOs:32, 34, 36, and 38 and the *Hordeum vulgare*, *Daucus carota*, and *Arabidopsis thaliana* sequences (SEQ ID NO:41, SEQ ID NO:42, and SEQ ID NO:43, respectively). The data in Table 8 represents a calculation of the percent identity of the amino acid sequences set forth in SEQ ID NOs:18, 32, 34, 36, and 38 and the *Hordeum vulgare*, *Daucus carota*, and *Arabidopsis thaliana* sequences (SEQ ID NO:41, SEQ ID NO:42, and SEQ ID NO:43, respectively)

TABLE 8

Percent Identity of Amino Acid Sequences Deduced From the Nucleotide Sequences of cDNA Clones Encoding Polypeptides Homologous to 4-Hydroxyphenylpyruvate Dioxygenase

SEQ ID NO.	Percent Identity to		
	3334222	3334219	3334223
18	49.4	58.2	60.0
32	58.5	73.3	67.4
34	86.4	73.3	69.9
36	49.4	58.2	60.0
38	94.2	59.9	58.1

Sequence alignments and percent identity calculations were performed using the Megalign program of the LASERGENE bioinformatics computing suite (DNASTAR Inc.,

Madison, WI). Multiple alignment of the sequences was performed using the Clustal method of alignment (Higgins and Sharp (1989) *CABIOS*. 5:151-153) with the default parameters (GAP PENALTY=10, GAP LENGTH PENALTY=10). Default parameters for pairwise alignments using the Clustal method were KTUPLE 1, GAP PENALTY=3,
5 WINDOW=5 and DIAGONALS SAVED=5. Sequence alignments and BLAST scores and probabilities indicate that the nucleic acid fragments comprising the instant cDNA clones encode substantial portions of rice, soybean, *Vernonia*, and wheat 4-hydroxyphenylpyruvate dioxygenase, and entire catalpa, soybean, and wheat 4-hydroxyphenylpyruvate dioxygenase. These sequences represent the first catalpa, rice, soybean, *Vernonia*, and
10 wheat sequences encoding 4-hydroxyphenylpyruvate dioxygenase.

EXAMPLE 5

Expression of Chimeric Genes in Monocot Cells

A chimeric gene comprising a cDNA encoding the instant polypeptides in sense orientation with respect to the maize 27 kD zein promoter that is located 5' to the cDNA fragment, and the 10 kD zein 3' end that is located 3' to the cDNA fragment, can be
15 constructed. The cDNA fragment of this gene may be generated by polymerase chain reaction (PCR) of the cDNA clone using appropriate oligonucleotide primers. Cloning sites (NcoI or SmaI) can be incorporated into the oligonucleotides to provide proper orientation of the DNA fragment when inserted into the digested vector pML103 as described below.
20 Amplification is then performed in a standard PCR. The amplified DNA is then digested with restriction enzymes NcoI and SmaI and fractionated on an agarose gel. The appropriate band can be isolated from the gel and combined with a 4.9 kb NcoI-SmaI fragment of the plasmid pML103. Plasmid pML103 has been deposited under the terms of the Budapest Treaty at ATCC (American Type Culture Collection, 10801 University Blvd., Manassas,
25 VA 20110-2209), and bears accession number ATCC 97366. The DNA segment from pML103 contains a 1.05 kb Sall-NcoI promoter fragment of the maize 27 kD zein gene and a 0.96 kb SmaI-Sall fragment from the 3' end of the maize 10 kD zein gene in the vector pGem9Zf(+) (Promega). Vector and insert DNA can be ligated at 15°C overnight, essentially as described (Maniatis). The ligated DNA may then be used to transform *E. coli*
30 XL1-Blue (Epicurian Coli XL-1 Blue™; Stratagene). Bacterial transformants can be screened by restriction enzyme digestion of plasmid DNA and limited nucleotide sequence analysis using the dideoxy chain termination method (Sequenase™ DNA Sequencing Kit; U.S. Biochemical). The resulting plasmid construct would comprise a chimeric gene encoding, in the 5' to 3' direction, the maize 27 kD zein promoter, a cDNA fragment
35 encoding the instant polypeptides, and the 10 kD zein 3' region.

The chimeric gene described above can then be introduced into corn cells by the following procedure. Immature corn embryos can be dissected from developing caryopses derived from crosses of the inbred corn lines H99 and LH132. The embryos are isolated 10

to 11 days after pollination when they are 1.0 to 1.5 mm long. The embryos are then placed with the axis-side facing down and in contact with agarose-solidified N6 medium (Chu et al. (1975) *Sci. Sin. Peking* 18:659-668). The embryos are kept in the dark at 27°C. Friable embryogenic callus consisting of undifferentiated masses of cells with somatic proembryoids and embryoids borne on suspensor structures proliferates from the scutellum of these immature embryos. The embryogenic callus isolated from the primary explant can be cultured on N6 medium and sub-cultured on this medium every 2 to 3 weeks.

The plasmid, p35S/Ac (obtained from Dr. Peter Eckes, Hoechst Ag, Frankfurt, Germany) may be used in transformation experiments in order to provide for a selectable marker. This plasmid contains the *Pat* gene (see European Patent Publication 0 242 236) which encodes phosphinothricin acetyl transferase (PAT). The enzyme PAT confers resistance to herbicidal glutamine synthetase inhibitors such as phosphinothricin. The *pat* gene in p35S/Ac is under the control of the 35S promoter from Cauliflower Mosaic Virus (Odell et al. (1985) *Nature* 313:810-812) and the 3' region of the nopaline synthase gene from the T-DNA of the Ti plasmid of *Agrobacterium tumefaciens*.

The particle bombardment method (Klein et al. (1987) *Nature* 327:70-73) may be used to transfer genes to the callus culture cells. According to this method, gold particles (1 µm in diameter) are coated with DNA using the following technique. Ten µg of plasmid DNAs are added to 50 µL of a suspension of gold particles (60 mg per mL). Calcium chloride (50 µL of a 2.5 M solution) and spermidine free base (20 µL of a 1.0 M solution) are added to the particles. The suspension is vortexed during the addition of these solutions. After 10 minutes, the tubes are briefly centrifuged (5 sec at 15,000 rpm) and the supernatant removed. The particles are resuspended in 200 µL of absolute ethanol, centrifuged again and the supernatant removed. The ethanol rinse is performed again and the particles resuspended in a final volume of 30 µL of ethanol. An aliquot (5 µL) of the DNA-coated gold particles can be placed in the center of a Kapton™ flying disc (Bio-Rad Labs). The particles are then accelerated into the corn tissue with a Biolistic™ PDS-1000/He (Bio-Rad Instruments, Hercules CA), using a helium pressure of 1000 psi, a gap distance of 0.5 cm and a flying distance of 1.0 cm.

For bombardment, the embryogenic tissue is placed on filter paper over agarose-solidified N6 medium. The tissue is arranged as a thin lawn and covered a circular area of about 5 cm in diameter. The petri dish containing the tissue can be placed in the chamber of the PDS-1000/He approximately 8 cm from the stopping screen. The air in the chamber is then evacuated to a vacuum of 28 inches of Hg. The macrocarrier is accelerated with a helium shock wave using a rupture membrane that bursts when the He pressure in the shock tube reaches 1000 psi.

Seven days after bombardment the tissue can be transferred to N6 medium that contains gluphosinate (2 mg per liter) and lacks casein or proline. The tissue continues to

grow slowly on this medium. After an additional 2 weeks the tissue can be transferred to fresh N6 medium containing glufosinate. After 6 weeks, areas of about 1 cm in diameter of actively growing callus can be identified on some of the plates containing the glufosinate-supplemented medium. These calli may continue to grow when sub-cultured on the

5 selective medium.

Plants can be regenerated from the transgenic callus by first transferring clusters of tissue to N6 medium supplemented with 0.2 mg per liter of 2,4-D. After two weeks the tissue can be transferred to regeneration medium (Fromm et al. (1990) *Bio/Technology* 8:833-839).

10

EXAMPLE 6

Expression of Chimeric Genes in Dicot Cells

A seed-specific expression cassette composed of the promoter and transcription terminator from the gene encoding the β subunit of the seed storage protein phaseolin from the bean *Phaseolus vulgaris* (Doyle et al. (1986) *J. Biol. Chem.* 261:9228-9238) can be used
15 for expression of the instant polypeptides in transformed soybean. The phaseolin cassette includes about 500 nucleotides upstream (5') from the translation initiation codon and about 1650 nucleotides downstream (3') from the translation stop codon of phaseolin. Between the 5' and 3' regions are the unique restriction endonuclease sites Nco I (which includes the ATG translation initiation codon), Sma I, Kpn I and Xba I. The entire cassette is flanked by
20 Hind III sites.

The cDNA fragment of this gene may be generated by polymerase chain reaction (PCR) of the cDNA clone using appropriate oligonucleotide primers. Cloning sites can be incorporated into the oligonucleotides to provide proper orientation of the DNA fragment when inserted into the expression vector. Amplification is then performed as described
25 above, and the isolated fragment is inserted into a pUC18 vector carrying the seed expression cassette.

Soybean embryos may then be transformed with the expression vector comprising sequences encoding the instant polypeptides. To induce somatic embryos, cotyledons, 3-5 mm in length dissected from surface sterilized, immature seeds of the soybean cultivar
30 A2872, can be cultured in the light or dark at 26°C on an appropriate agar medium for 6-10 weeks. Somatic embryos which produce secondary embryos are then excised and placed into a suitable liquid medium. After repeated selection for clusters of somatic embryos which multiplied as early, globular staged embryos, the suspensions are maintained as described below.

35 Soybean embryogenic suspension cultures can be maintained in 35 mL liquid media on a rotary shaker, 150 rpm, at 26°C with florescent lights on a 16:8 hour day/night schedule. Cultures are subcultured every two weeks by inoculating approximately 35 mg of tissue into 35 mL of liquid medium.

Soybean embryogenic suspension cultures may then be transformed by the method of particle gun bombardment (Klein et al. (1987) *Nature* (London) 327:70-73, U.S. Patent No. 4,945,050). A DuPont Biolistic™ PDS1000/HE instrument (helium retrofit) can be used for these transformations.

- 5 A selectable marker gene which can be used to facilitate soybean transformation is a chimeric gene composed of the 35S promoter from Cauliflower Mosaic Virus (Odell et al. (1985) *Nature* 313:810-812), the hygromycin phosphotransferase gene from plasmid pJR225 (from *E. coli*; Gritz et al. (1983) *Gene* 25:179-188) and the 3' region of the nopaline synthase gene from the T-DNA of the Ti plasmid of *Agrobacterium tumefaciens*. The seed expression
10 cassette comprising the phaseolin 5' region, the fragment encoding the instant polypeptides and the phaseolin 3' region can be isolated as a restriction fragment. This fragment can then be inserted into a unique restriction site of the vector carrying the marker gene.

- To 50 μ L of a 60 mg/mL 1 μ m gold particle suspension is added (in order): 5 μ L DNA (1 μ g/ μ L), 20 μ l spermidine (0.1 M), and 50 μ L CaCl_2 (2.5 M). The particle
15 preparation is then agitated for three minutes, spun in a microfuge for 10 seconds and the supernatant removed. The DNA-coated particles are then washed once in 400 μ L 70% ethanol and resuspended in 40 μ L of anhydrous ethanol. The DNA/particle suspension can be sonicated three times for one second each. Five μ L of the DNA-coated gold particles are then loaded on each macro carrier disk.

- 20 Approximately 300-400 mg of a two-week-old suspension culture is placed in an empty 60x15 mm petri dish and the residual liquid removed from the tissue with a pipette. For each transformation experiment, approximately 5-10 plates of tissue are normally bombarded. Membrane rupture pressure is set at 1100 psi and the chamber is evacuated to a vacuum of 28 inches mercury. The tissue is placed approximately 3.5 inches away from the
25 retaining screen and bombarded three times. Following bombardment, the tissue can be divided in half and placed back into liquid and cultured as described above.

- Five to seven days post bombardment, the liquid media may be exchanged with fresh media, and eleven to twelve days post bombardment with fresh media containing 50 mg/mL hygromycin. This selective media can be refreshed weekly. Seven to eight weeks post
30 bombardment, green, transformed tissue may be observed growing from untransformed, necrotic embryogenic clusters. Isolated green tissue is removed and inoculated into individual flasks to generate new, clonally propagated, transformed embryogenic suspension cultures. Each new line may be treated as an independent transformation event. These suspensions can then be subcultured and maintained as clusters of immature embryos or
35 regenerated into whole plants by maturation and germination of individual somatic embryos.

EXAMPLE 7

Expression of Chimeric Genes in Microbial Cells

The cDNAs encoding the instant polypeptides can be inserted into the T7 *E. coli* expression vector pBT430. This vector is a derivative of pET-3a (Rosenberg et al. (1987) *Gene* 56:125-135) which employs the bacteriophage T7 RNA polymerase/T7 promoter system. Plasmid pBT430 was constructed by first destroying the EcoR I and Hind III sites in pET-3a at their original positions. An oligonucleotide adaptor containing EcoR I and Hind III sites was inserted at the BamH I site of pET-3a. This created pET-3aM with additional unique cloning sites for insertion of genes into the expression vector. Then, the Nde I site at the position of translation initiation was converted to an Nco I site using oligonucleotide-directed mutagenesis: The DNA sequence of pET-3aM in this region, 5'-CATATGG, was converted to 5'-CCCATGG in pBT430.

Plasmid DNA containing a cDNA may be appropriately digested to release a nucleic acid fragment encoding the protein. This fragment may then be purified on a 1% NuSieve GTG™ low melting agarose gel (FMC). Buffer and agarose contain 10 µg/ml ethidium bromide for visualization of the DNA fragment. The fragment can then be purified from the agarose gel by digestion with GELase™ (Epicentre Technologies) according to the manufacturer's instructions, ethanol precipitated, dried and resuspended in 20 µL of water. Appropriate oligonucleotide adapters may be ligated to the fragment using T4 DNA ligase (New England Biolabs, Beverly, MA). The fragment containing the ligated adapters can be purified from the excess adapters using low melting agarose as described above. The vector pBT430 is digested, dephosphorylated with alkaline phosphatase (NEB) and deproteinized with phenol/chloroform as described above. The prepared vector pBT430 and fragment can then be ligated at 16°C for 15 hours followed by transformation into DH5 electrocompetent cells (GIBCO BRL). Transformants can be selected on agar plates containing LB media and 100 µg/mL ampicillin. Transformants containing the gene encoding the instant polypeptides are then screened for the correct orientation with respect to the T7 promoter by restriction enzyme analysis.

For high level expression, a plasmid clone with the cDNA insert in the correct orientation relative to the T7 promoter can be transformed into *E. coli* strain BL21(DE3) (Studier et al. (1986) *J. Mol. Biol.* 189:113-130). Cultures are grown in LB medium containing ampicillin (100 mg/L) at 25°C. At an optical density at 600 nm of approximately 1, IPTG (isopropylthio-β-galactoside, the inducer) can be added to a final concentration of 0.4 mM and incubation can be continued for 3 h at 25°. Cells are then harvested by centrifugation and re-suspended in 50 µL of 50 mM Tris-HCl at pH 8.0 containing 0.1 mM DTT and 0.2 mM phenyl methylsulfonyl fluoride. A small amount of 1 mm glass beads can be added and the mixture sonicated 3 times for about 5 seconds each time with a microprobe sonicator. The mixture is centrifuged and the protein concentration of the supernatant

determined. One μg of protein from the soluble fraction of the culture can be separated by SDS-polyacrylamide gel electrophoresis. Gels can be observed for protein bands migrating at the expected molecular weight.

EXAMPLE 8

5 Evaluating Compounds for Their Ability to Inhibit the Activity of Vitamin E Biosynthetic Enzymes

10 The polypeptides described herein may be produced using any number of methods known to those skilled in the art. Such methods include, but are not limited to, expression in bacteria as described in Example 7, or expression in eukaryotic cell culture, *in planta*, and using viral expression systems in suitably infected organisms or cell lines. The instant polypeptides may be expressed either as mature forms of the proteins as observed *in vivo* or as fusion proteins by covalent attachment to a variety of enzymes, proteins or affinity tags. Common fusion protein partners include glutathione S-transferase ("GST"), thioredoxin ("Trx"), maltose binding protein, and C- and/or N-terminal hexahistidine polypeptide ("His₆"). The fusion proteins may be engineered with a protease recognition site at the fusion point so that fusion partners can be separated by protease digestion to yield intact mature enzyme. Examples of such proteases include thrombin, enterokinase and factor Xa. However, any protease can be used which specifically cleaves the peptide connecting the fusion protein and the enzyme.

20 Purification of the instant polypeptides, if desired, may utilize any number of separation technologies familiar to those skilled in the art of protein purification. Examples of such methods include, but are not limited to, homogenization, filtration, centrifugation, heat denaturation, ammonium sulfate precipitation, desalting, pH precipitation, ion exchange chromatography, hydrophobic interaction chromatography and affinity chromatography, wherein the affinity ligand represents a substrate, substrate analog or inhibitor. When the instant polypeptides are expressed as fusion proteins, the purification protocol may include the use of an affinity resin which is specific for the fusion protein tag attached to the expressed enzyme or an affinity resin containing ligands which are specific for the enzyme. For example, the instant polypeptides may be expressed as a fusion protein coupled to the C-terminus of thioredoxin. In addition, a (His)₆ peptide may be engineered into the N-terminus of the fused thioredoxin moiety to afford additional opportunities for affinity purification. Other suitable affinity resins could be synthesized by linking the appropriate ligands to any suitable resin such as Sepharose-4B. In an alternate embodiment, a thioredoxin fusion protein may be eluted using dithiothreitol; however, elution may be accomplished using other reagents which interact to displace the thioredoxin from the resin. These reagents include β -mercaptoethanol or other reduced thiol. The eluted fusion protein may be subjected to further purification by traditional means as stated above, if desired. Proteolytic cleavage of the thioredoxin fusion protein and the enzyme may be accomplished

after the fusion protein is purified or while the protein is still bound to the ThioBond™ affinity resin or other resin.

Crude, partially purified or purified enzyme, either alone or as a fusion protein, may be utilized in assays for the evaluation of compounds for their ability to inhibit enzymatic activation of the instant polypeptides disclosed herein. Assays may be conducted under well known experimental conditions which permit optimal enzymatic activity. For example, assays for gamma tocopherol methyltransferase are presented by d'Harlingue and Camara (1985) *J. Biol. Chem.* 260:15200-15203. Assays for 4-hydroxyphenylpyruvate dioxygenase are presented by Norris et al. (1995) *Plant Cell* 7: 2139-2149.

10

EXAMPLE 9

Expression of gamma Tocopherol Methyltransferase in Soybean Somatic Embryos

The ability to change the levels of alpha- and gamma-tocopherol in plants by transforming them with sequences encoding gamma tocopherol methyltransferase was tested by preparing transgenic soybean somatic embryos and assaying the tocopherol levels. Plasmid DNA from clone sah1c.pk001.k8 was used as a template to prepare a Not I pcr fragment encoding the entire deduced open reading frame using the following pcr primers (forward primer AGC GCG GCC GCA TGG CCA CCG TGG TGA GGA TCC CAA CAA TCT CAT GCA TCC ACA TCC ACA; reverse primer AGC GCG GCC GCT TAT CTA GTG TGG AAA TAA TGA TCA). Standard pcr reactions were used (100 microliter total reaction containing 5 ng plasmid, 25 nmoles primer, 25 nmoles dNTPs, 1x cloned pfu buffer (Stratagene), 5% DMSO, 5U cloned pfu DNA polymerase (Stratagene). The recommended cycling parameters for pcr with pfu DNA polymerase were used (denature 45 s 94°C, anneal 45 s 55°C, extend 2.5 min 72°C, 25 cycles, final 10min extension at 72°C). The pcr product was purified on a 1% agarose/TAE gel (precast FMC), the ethidium bromide visualized band cut out and purified using a QIAquick gel extraction kit (Qiagen). The band prep (40 ng) was ligated into PCR Blunt (InVitrogen) according to the manufacturers recommendations, and the ligated plasmid used to transform *E. coli* DH10 cells. Kanamycin resistant colonies were grown overnight in liquid culture (LB/Kan), plasmids prepared and cut with Not I. Plasmids containing the correct insert size were selected for full insert sequence to confirm fidelity of pcr. Plasmids containing the correct insert verified by DNA sequencing were digested with Not I and ligated to Not I-digested and phosphatase-treated pKS67. The plasmid pKS67 was prepared by replacing in pRB20 (described in U.S. Patent No. 5,846,784) the 800 bp Nos 3' fragment, with the 285 bp Nos 3' fragment containing the polyadenylation signal sequence and described in Depicker et al. (1982) *J. Mol. Appl. Genet.* 1:561-573. Clones were screened for the sense and anti-sense orientation of the gamma tocopherol methyltransferase insert fragment by restriction enzyme digestion.

Transformation of Soybean Somatic Embryo Cultures

The following stock solutions and media were used for transformation and propagation of soybean somatic embryos:

Stock Solutions		Media
<u>MS Sulfate 100x stock</u> (g/L)		<u>SB55 (per Liter)</u>
MgSO ₄ ·7H ₂ O	37.0	10 mL of each MS stock
MnSO ₄ ·H ₂ O	1.69	1 mL of B5 Vitamin stock
ZnSO ₄ ·7H ₂ O	0.86	0.8 g NH ₄ NO ₃
CuSO ₄ ·5H ₂ O	0.0025	3.033 g KNO ₃
		1 mL 2,4-D (10 mg/mL stock)
		0.667 g asparagine
		pH 5.7
<u>MS Halides 100x stock</u>		<u>SB103 (per Liter)</u>
CaCl ₂ ·2H ₂ O	44.0	1 pk. Murashige & Skoog salt mixture*
KI	0.083	60 g maltose
CoCl ₂ ·6H ₂ O	0.00125	2 g gelrite
KH ₂ PO ₄	17.0	pH 5.7
H ₃ BO ₃	0.62	
Na ₂ MoO ₄ ·2H ₂ O	0.025	
Na ₂ EDTA	3.724	
FeSO ₄ ·7H ₂ O	2.784	
		<u>SB148 (per Liter)</u>
<u>B5 Vitamin stock</u>		1 pk. Murashige & Skoog salt mixture*
myo-inositol	100.0	60 g maltose
nicotinic acid	1.0	1 mL B5 vitamin stock
pyridoxine HCl	1.0	7 g agarose
thiamine	10.0	pH 5.7
		*(Gibco BRL)

5

Soybean embryonic suspension cultures were maintained in 35 mL liquid media (SB55) on a rotary shaker (150 rpm) at 28°C with a mix of fluorescent and incandescent lights providing a 16 h day 8 h night cycle. Cultures were subcultured every 2 to 3 weeks by inoculating approximately 35 mg of tissue into 35 mL of fresh liquid media.

10

Soybean embryonic suspension cultures were transformed with the plasmid containing the gamma tocopherol methyltransferase sequence by the method of particle gun bombardment (see Klein et al. (1987) *Nature* 327:70-73) using a DuPont Biolistic PDS1000/He instrument. Five µL of pKS93s plasmid DNA (1 g/L), 50 µL CaCl₂ (2.5 M), and 20 µL spermidine (0.1 M) were added to 50 µL of a 60 mg/mL 1 mm gold particle suspension. The particle preparation was agitated for 3 minutes, spun on a microfuge for

15

10 seconds and the supernate removed. The DNA-coated particles were then washed once with 400 μ L of 70% ethanol and resuspended in 40 μ L of anhydrous ethanol. The DNA/particle suspension was sonicated three times for 1 second each. Five μ L of the DNA-coated gold particles were then loaded on each macro carrier disk.

- 5 Approximately 300 to 400 mg of two-week-old suspension culture was placed in an empty 60 mm X 15 mm petri dish and the residual liquid removed from the tissue using a pipette. The tissue was placed about 3.5 inches away from the retaining screen and bombarded twice. Membrane rupture pressure was set at 1100 psi and the chamber was evacuated to -28 inches of Hg. Two plates were bombarded, and following bombardment, 10 the tissue was divided in half, placed back into liquid media, and cultured as described above.

- Fifteen days after bombardment, the liquid media was exchanged with fresh SB55 containing 50 mg/mL hygromycin. The selective media was refreshed weekly. Six weeks after bombardment, green, transformed tissue was isolated and inoculated into flasks to 15 generate new transformed embryonic suspension cultures.

- Transformed embryonic clusters were removed from liquid culture media and placed on a solid agar media, SB103, containing 0.5% charcoal to begin maturation. After 1 week, embryos were transferred to SB103 media minus charcoal. After 5 weeks on SB103 media, maturing embryos were separated and placed onto SB148 media. During maturation 20 embryos were kept at 26°C with a mix of fluorescent and incandescent lights providing a 16 h day 8 h night cycle. After 3 weeks on SB148 media, embryos were analyzed for the expression of the tocopherols. Each embryonic cluster gave rise to 5 to 20 somatic embryos.

 Non-transformed somatic embryos were cultured by the same method as used for the transformed somatic embryos.

25 *Analysis of Transformed Somatic Embryos*

- At the end of the 6th week on SB148 medium somatic embryos were harvested from 25 independently transformed lines. Somatic embryos were collected in pools of five and weighed for fresh weight. Excess embryos were stored in 96-well plates at -80°. The pooled somatic embryos were lyophilized for 18 hours and the dry weight measured. The 30 lyophilized somatic embryos were briefly pulverised with a hand held Potter homogeniser and then 600 μ L of heptane added and the samples incubated for 24 hours in the dark at room temperature to extract oils and tocopherols. The heptane was decanted and a further 300 μ L added to the samples. The extracts were combined and centrifuged (5 min, 12000 g). The supernatant was stored in amber hplc autosampler vials at -20°C prior to analysis.

- 35 HPLC analysis of the extracts was carried out using an HP1100 system (Agilent Technologies) 25 μ L of the heptane sample was applied to a Lichrosphere Si 60 column (5 micron 4 x 12.5 mm). The column was eluted with heptane/isopropanol (98:2 v/v) at a flow rate of 1 mL/min. After 6 minutes all four tocopherol isomers were eluted, as detected by

a HP1100 fluorescence detector (Excitation wavelength 295 nm, emission wavelength 330 nm). Individual tocopherol standards (Matreya) were diluted with hplc grade heptane to levels between 1 and 200 ng/ul to construct a 6 point external standard curve. Tocopherols in each oil were quantified using a standard curve run on the same day as the samples. The

5 sum of tocopherol peak areas of samples from a non-transformed control line were compared with those of 25 independent gamma tocopherol methyltransferase-transformed, hygromycin resistant lines. Figure 3 shows a graph depicting the distribution of the percent alpha- and gamma-tocopherol of soybean somatic embryos transgenic for the gamma tocopherol methyltransferase gene and a control line. The mean alpha tocopherol content is 40 to 70%.

10 Some lines, such as the ones represented in bars 2, 5, and 18 appear to represent overexpression of the gamma tocopherol methyltransferase, which would be predicted to yield higher levels of alpha tocopherol. Other lines, such as those represented in bars 6 and 10 appear to represent co-suppression of the gamma tocopherol methyltransferase since this is predicted to yield higher levels of gamma tocopherol. These results indicate that

15 transgenic expression of gamma-tocopherol methyltransferase affords the ability to manipulate tocopherol levels as desired for a particular application.

Various modifications of the invention in addition to those shown and described herein will be apparent to those skilled in the art from the foregoing description. Such modifications are also intended to fall within the scope of the appended claims.

20 The disclosure of each reference set forth above is incorporated herein by reference in its entirety.

CLAIMS

What is claimed is:

1. An isolated polynucleotide comprising a first nucleotide sequence encoding a polypeptide of at least 160 amino acids that has at least 80% identity based on the Clustal method of alignment when compared to a polypeptide selected from the group consisting of SEQ ID NOs:2, 4, 6, 8, 10, 22, 24, 26, 28, and 30, or a second nucleotide sequence comprising the complement of the first nucleotide sequence.
2. An isolated polynucleotide comprising a first nucleotide sequence encoding a polypeptide of at least 150 amino acids that has at least 95% identity based on the Clustal method of alignment when compared to a polypeptide selected from the group consisting of SEQ ID NOs:12, 14, 16, 18, 20, 32, 34, 36, and 38, or a second nucleotide sequence comprising the complement of the first nucleotide sequence.
3. The isolated polynucleotide of Claim 1 or Claim 2, wherein the first nucleotide sequence consists of a nucleic acid sequence selected from the group consisting of SEQ ID NOs:1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, and 37 that codes for the polypeptide selected from the group consisting of SEQ ID NOs:2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, and 38.
4. The isolated polynucleotide of Claim 1 or Claim 2 wherein the nucleotide sequences are DNA.
5. The isolated polynucleotide of Claim 1 or Claim 2 wherein the nucleotide sequences are RNA.
6. A chimeric gene comprising the isolated polynucleotide of Claim 1 or Claim 2 operably linked to suitable regulatory sequences.
7. An isolated host cell comprising the chimeric gene of Claim 6.
8. A host cell comprising an isolated polynucleotide of Claim 1 or Claim 2.
9. The host cell of Claim 8 wherein the host cell is selected from the group consisting of yeast, bacteria, plant, and virus.
10. A virus comprising the isolated polynucleotide of Claim 1 or Claim 2.
11. A polypeptide of at least 160 amino acids that has at least 80% identity based on the Clustal method of alignment when compared to a polypeptide selected from the group consisting of SEQ ID NOs:2, 4, 6, 8, 10, 22, 24, 26, 28, and 30.
12. A polypeptide of at least 150 amino acids that has at least 95% identity based on the Clustal method of alignment when compared to a polypeptide selected from the group consisting of SEQ ID NOs:12, 14, 16, 18, 20, 32, 34, 36, and 38.
13. A method of selecting an isolated polynucleotide that affects the level of expression of a vitamin E biosynthetic enzyme polypeptide in a plant cell, the method comprising the steps of:

(a) constructing an isolated polynucleotide comprising a nucleotide sequence of at least one of 30 contiguous nucleotides derived from an isolated polynucleotide of Claim 1 or Claim 2;

(b) introducing the isolated polynucleotide into a plant cell; and

5 (c) measuring the level of a polypeptide in the plant cell containing the polynucleotide to provide a positive selection means.

14. The method of Claim 13 wherein the isolated polynucleotide consists of a nucleotide sequence selected from the group consisting of SEQ ID NOs:1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, and 37 that codes for the polypeptide selected from
10 the group consisting of SEQ ID NOs:2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, and 38.

15 15. A method of selecting an isolated polynucleotide that affects the level of expression of a vitamin E biosynthetic enzyme polypeptide in a plant cell, the method comprising the steps of:

(a) constructing an isolated polynucleotide of Claim 1 or Claim 2;

(b) introducing the isolated polynucleotide into a plant cell; and

(c) measuring the level of polypeptide in the plant cell containing the polynucleotide to provide a positive selection means.

16. A method of obtaining a nucleic acid fragment encoding a vitamin E
20 biosynthetic enzyme polypeptide comprising the steps of:

(a) synthesizing an oligonucleotide primer comprising a nucleotide sequence of at least one of 30 contiguous nucleotides derived from a nucleotide sequence selected from the group consisting of SEQ ID NOs:1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, and the complement of such nucleotide sequences; and

25 (b) amplifying a nucleic acid sequence using the oligonucleotide primer.

17. A method of obtaining a nucleic acid fragment encoding a vitamin E biosynthetic enzyme polypeptide comprising the steps of:

(a) probing a cDNA or genomic library with an isolated polynucleotide comprising at least one of 30 contiguous nucleotides derived from a nucleotide sequence
30 selected from the group consisting of SEQ ID NOs:1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37 and the complement of such nucleotide sequences;

(b) identifying a DNA clone that hybridizes with the isolated polynucleotide;

(c) isolating the identified DNA clone; and

(d) sequencing the cDNA or genomic fragment that comprises the isolated DNA
35 clone.

18. A method for evaluating at least one compound for its ability to inhibit the activity of a vitamin E biosynthetic enzyme, the method comprising the steps of:

(a) transforming a host cell with a chimeric gene comprising a nucleic acid fragment encoding a vitamin E biosynthetic enzyme, operably linked to suitable regulatory sequences;

5 (b) growing the transformed host cell under conditions that are suitable for expression of the chimeric gene wherein expression of the chimeric gene results in production of the vitamin E biosynthetic enzyme encoded by the operably linked nucleic acid fragment in the transformed host cell;

(c) optionally purifying the vitamin E biosynthetic enzyme expressed by the transformed host cell;

10 (d) treating the vitamin E biosynthetic enzyme with a compound to be tested; and

(e) determining the activity of the vitamin E biosynthetic enzyme that has been treated with the compound, thereby selecting compounds with potential for inhibitory activity.

15 19. A composition comprising the isolated polynucleotide of Claim 1 or Claim 2.

20. A composition comprising the isolated polypeptide of Claim 11.

21. An isolated polynucleotide comprising the nucleotide sequence having at least one of 30 contiguous nucleotides derived from a nucleic acid sequence selected from the group consisting of SEQ ID NOs: 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33,
20 35, 37 and the complement of such sequences.

22. An expression cassette comprising an isolated polynucleotide of Claim 1 or Claim 2 operably linked to a promoter.

23. A method for positive selection of a transformed cell comprising:

25 (a) transforming a host cell with an expression cassette of Claim 22; and
(b) growing the transformed host cell under conditions which allow expression of the polynucleotide in an amount sufficient to complement a null mutant to provide a positive selection means.

24. The method of Claim 23 wherein the plant cell is a monocot.

25. The method of Claim 23 wherein the plant cell is a dicot.

30

FIGURE 1

SEQ ID NO: 39	MVYHVRPKHAL-----FLAFYCYFS-----	60
SEQ ID NO: 40	MKATLAAPSSLTSL-P-----YRTNSS-FGSKSSLLFRSPSSSSSVS-MTTTRGNV	
SEQ ID NO: 38	MAN-----SXRPA-PLTPLHRLDAAPPRLSLGHAARPVPREVLPPLPAR-XLRAPDGV	
SEQ ID NO: 28	MATVVRI-----PTISCIHIHTRFSQSPRTFARIRVGPRSWAPIRASA-ASSERGEI	
SEQ ID NO: 08	MATVVRI-----PTISCIHIHTRFSQSPRTFARIRVGPRSWAPIRASA-ASSERGEI	
SEQ ID NO: 22	MAHAALLHCSQSSR-SLAACRRGSHYRAPSH-VPRHSRRLRAVVSL---R-PMASSTA-	
SEQ ID NO: 24	AR--VQPTGALAPLHPLLRCTSRHLCASAPRAGLCLEHHRRRRSSRRRTKLAVRAMAPT	
SEQ ID NO: 26	FRH-----GH-----AL-----	
	1	
SEQ ID NO: 39	LLTMASATIASADLYEKIKNEFYDDSSGLWEDVWGEHMHGYYGPHGTYRI---DRRQAQI	
SEQ ID NO: 40	AVAAAATSTEALRKGAIEFYNETSGLWEEIWGDHMHGFDYDPDSSVQLSDSGHKEAQI	
SEQ ID NO: 38	VDDRGPGDAAPPGLKEGIAGLYDESSGLWESIWEHMHGFDYDSGEAASMSD--HRAAQI	
SEQ ID NO: 28	VLEQPKKDDKKLQKQIAEFYDESSGLWENIWGDHMHGFDYDSDTVSLSD--HRAAQI	
SEQ ID NO: 08	VLEQPKKDDKKLQKQIAEFYDESSGLWENIWGDHMHGFDYDSDTVSLSD--HRAAQI	
SEQ ID NO: 22	---QAPATAPPGLKEGIAGLYDESSGLWENIWGDHMHGFDYDSEAAASMD--HRAAQI	
SEQ ID NO: 24	LSSSSTAAPGLKEGIAGLYDESSGVWESIWEHMHGFDYDAGEAASMSD--HRAAQI	
SEQ ID NO: 26	-----	
	61	
SEQ ID NO: 39	DLIKELLAWAVPQNSA---KPRKILDGCGIGGSSLYLAQQHQAQAEVMGASLSPVQVERA	120
SEQ ID NO: 40	RMIEESLRFAGVTDEEE-EKKIKKVVDVCGGIGGSSRYLASKFGAECIGITLSPVQAKRA	
SEQ ID NO: 38	RMIEEALAFAAVP--DDPTNPKPTIVDVGCGIGGSSRYLANKYGAQCSTLSPVQAERG	
SEQ ID NO: 28	RMIQESLRFASV--SEERSKWPKSIVDVGCGIGGSSRYLAKKFGATSVGITLSPVQAQRA	
SEQ ID NO: 08	RMIQESLRFASV--SEERSKWPKSIVDVGCGIGGSSRYLAKKFGATSVGITLSPVQAQRA	
SEQ ID NO: 22	RMIEEALAFAGVPASDDPEKTPKTIIVDVGCGIGGSSRYLAKKYGXQCTGITLSPVQAERG	
SEQ ID NO: 24	RMIEESLAF-----	
SEQ ID NO: 26	-----	
	121	
SEQ ID NO: 39	-----	180

FIGURE 1

```

SEQ ID NO:39 GERARALGLSTCQFQVANALDLPFASDSFDWVWSLESGEHMPNKAQFLQEAWRVLKPGG
SEQ ID NO:40 NDIAAQSLSHKASFQVADALDQPFEDGKFDLVWSMESGEHMPDKAKFVKELVRVAAPGG
SEQ ID NO:38 NALAAAQGLSDKASFQVADALEQFPDGGFDLVWSMESGEHMPNKKQFVSELARVAAPGA
SEQ ID NO:28 NALAAAQGLADKVSFQVADALQQPFSDGGFDLVWSMESGEHMPDKAKFVGELARVAAPGA
SEQ ID NO:08 NALAAAQGLADKVSFQVADALQQPFSDGGFDLVWSMESGEHMPDKAKFVGELARVAAPGA
SEQ ID NO:22 NALAAAQGLSDQVTLQVADALEQFPDGGFDLVWSMESGEHMPDKRKFVSELARVAAPGG
SEQ ID NO:24 -----
SEQ ID NO:26 -----AQFPDGGFDLVWSMESDEHMPDKRQFVSELARVAAPGA
181 240

SEQ ID NO:39 RLILATWCHRPIDPGNGPLTADERRHLQAIYDVYCLPYVVSLLPDYEAIARECGFGEIKTA
SEQ ID NO:40 RIIIVTWCHRNLSAGEEALQPWEQNILDKICKTFYLPAWCSTDDYVNLQSHSLQDIKCA
SEQ ID NO:38 TIIIVTWCHRNLPSEDSLKPDELNLKKICDAYYLPDWCSPSDYVKIAESLSLEDIKTA
SEQ ID NO:28 IIIIVTWCHRDLPDEQSLHPWEQDLLKKICDAYYLPAWCSTSDYVKLLQSLSLQDIKSE
SEQ ID NO:08 IIIIVTWCHRDLPDEQSLHPWEQDLLKKICDAYYLPAWCSTSDYVKLLQSLSLQDIKSE
SEQ ID NO:22 TIIIVTWCHRNLDPSETSLKPDELSLLRRICDAYYLPDWCSPSDYVNIAKSLSLEDIKTA
SEQ ID NO:24 -----
SEQ ID NO:26 RIIIVTWCHRNLEPSEESLKPDELNLKKRICDAYYLPDWCSPSDYVKIAESLSLEDIRTA
241 300

SEQ ID NO:39 DWSVAVAPFWDRVIESAFDPRLWALGQAGPKIINAALCLRLMKWYERGLVRFGLLTGI
SEQ ID NO:40 DWSENVAPFWPAPVIRTALTWKGLVSLRSGMKS IKGALTMPLMIEGYKKGVIKFGIITCQ
SEQ ID NO:38 DWSENVAPFWPAPVIRSAFTWKGLTSLRSGWKT IKGALVMPMLIQGYKKGLIKFSIITCR
SEQ ID NO:28 DWSRFVAPFWPAPVIRSAFTWKGLSLLSSGQKT IKGALAMPLMIEGYKKDLIKFAIITCR
SEQ ID NO:08 DWSRFVAPFWPAPVIRSAFTWKGLSLLSSGKLG I-----YIAFQKQTPPSSIATCK
SEQ ID NO:22 DWSENVAPFWPAPVIRSAFTWKGLTSLTTGWKTIRGAMVMPMLIQGYKKGLIKFTIITCR
SEQ ID NO:24 -----
SEQ ID NO:26 DWSENVAPFWPAPVIRSAFTWKGLTSLRSGWETVRGAMVMPMLVIEGYKKGLIKFPIITCR
301 360

```

FIGURE 1

SEQ ID NO:39	KPLV-----	361	380
SEQ ID NO:40	KPL-----		
SEQ ID NO:38	KPQAAIEGEPEAAPSVE.-		
SEQ ID NO:28	K-----PE.-		
SEQ ID NO:08	SYVTDHYFHTR.-		
SEQ ID NO:22	KPGAA-.MA---HAALLHC		
SEQ ID NO:24	-----		
SEQ ID NO:26	KPETT-----Q.-		

FIGURE 2

SEQ ID NO: 41	MPPTPTT PAATGAAA AVTPEHARPH-----RMVRFNPRSDREHTLSFHHVEFWCADAASA	60
SEQ ID NO: 43	MGHQNAAVSENQNHDDGAASSPGFKLVGFSKFVRKNPKSKDKFKVKREHHIEFWCGDATNV	
SEQ ID NO: 42	MGKKQSEAEILSS-NSSNTSPATFKLVGFNFVRANPKSDHFAVKREHHIEFWCGDATNT	
SEQ ID NO: 36	MPIPMCNEIQQA-QAQAQAQPGFKLVGFKNFVRTNPKSDREQVNRHHIEFWCTDATNA	
SEQ ID NO: 38	MPPTPTT PAATGAAA-VTPEHARPR-----RMVRFNPRSDREHTLAFHHVEFWCADAASA	
SEQ ID NO: 32	MGK-QTTTSATAA-DGSKDAHAEEFKLVGFKNFVRTNPKSDHFCVHRHHIEFWCGDATNT	1
		60
SEQ ID NO: 41	AGRFALGAPLAARSDLSTGNSAHASQLLRSGSLAFLFTAPYANG-----CDAATASL	
SEQ ID NO: 43	ARRFSWGLGMRFSAKSDLSTGNMVMHASYLITSGDLREFTAPYSPSLSAGEIKPTTTASI	
SEQ ID NO: 42	SRRFSWGLGMPILVAKSDLSTGNSVHASYLVRSANLSEVFTAPYSPSTTTSS-G---SAAI	
SEQ ID NO: 36	SRRFSWGLGMPILVAKSDLSTGNIHASYLRLSGDLSEFLSAPYSPSLSAGS-SAASSASI	
SEQ ID NO: 38	AGRFALGAPLAARSDLSTGNSVHASQLLRSGNLAFLEFTAPYANG-----CDAATASL	
SEQ ID NO: 32	AKRFSWGLGMPILVAKSDLSTGNSAHASYLRLSGELNFLTSPYSPSISAPS-----SAAI	61
		120
SEQ ID NO: 41	PSFSADAARRFSADHGI AVRSV ALRVADAAEA FRASRRRGARPAFAPVDLGRGEA-FAEV	
SEQ ID NO: 43	PSFDHGSCRSFFSSHGLGVRAVAIEVEDAESAFSISVANGAIPSSPPIVLNEAVT-IAEV	
SEQ ID NO: 42	PSFSASGFHSFAAKHGLAVRAIALEVADVAFAAFASVARGARPA SAPVEL-DDQAWLAEV	
SEQ ID NO: 36	PSFDAATCLAFAAKHGFGVRAIALEVADAAEAFAFSVAKGAEPASPPV-LVDDRTGFAEV	
SEQ ID NO: 38	PSFSADAARQFSADHGLAVRSIALRVADAAEA FRASVDGGARPAFSPVDLGRGFG-FAEV	
SEQ ID NO: 32	PSFSFSTYQSFTSSHGLAVRAVAIQVDSAFSAFSAYSASISRGAKPV SAPILLSDNKTAIAEV	121
		180

FIGURE 2

```

SEQ ID NO:41  ELYGDIVLRFVSH-PDGTDP-----FLPGFEGVTNPD-----VDYGLTRFDHVVGNVP
SEQ ID NO:43  KLYGDIVLRYVSYKAEDTEKSE-----FLPGFERVEDASSFP-LDYGIRRLDHAVGNVP
SEQ ID NO:42  ELYGDIVLRFVSFGRE-----EGLFLPGFEAVEGTASFDPDLDYGIRRLDHAVGNVT
SEQ ID NO:36  RLYGDIVLRYVSYKDAAPQAPHADPSRWFLPGFEAAAASSSFPELDYGIRRLDHAVGNVP
SEQ ID NO:38  ELYGDIVLRFVSH-PDGRDVP-----FLPGFEGVSNPD-----VDYGLTRFDHVVGNVP
SEQ ID NO:32  HLYGDSVLRVFSYGDNG-----TGPDGWFLPGFEPVDDQMSYKELDYGIRRLDHAVGNVP
                                                    181      240

SEQ ID NO:41  ELAPAAAYIAGFTGFHEFAEFTAEDVGTTESGLNSVVLANNSEGVLLPLNEPVHGTKRRS
SEQ ID NO:43  ELGPALTYVAGFTGFHQFAEFTADDVGTAEGLNSAVLASNDEMVLPLINEPVHGTKRKS
SEQ ID NO:42  ELGPVVEYIKGFTGFHEFAEFTAEDVGTLESGLNSVVLANNSEMVLLPLNEPVYGTKRKS
SEQ ID NO:36  ELAPAVRYLKGFSGFHEFAEFTAEDVGTSEGLNSVVLANNSETVLLPLNEPVYGTKRKS
SEQ ID NO:38  ELAPAAAYVAGFTGFHEFAEFTTEDVGTAEGLNSMVLANNSEGVLLPLNEPVHGTKRRS
SEQ ID NO:32  ELGPVVDYLLKKFTGFHEFAEFTSEDVGTAEGLNSMVLANNENVLLPLNEPVFGTKRKS
                                                    241      300


SEQ ID NO:41  QIQTLEHHGGPGVQHI AVASSDVLRTLRKMRRARSAMGGDFLPPPLPKYYEGVRRLAGD
SEQ ID NO:43  QIQTYLEHNEGAGLQHLALMSEDI FRTLREMRKRSSIGGDFMPSPPTYYQNLKRVGD
SEQ ID NO:42  QIQTYLEHNEGAGVQHLALVSEDI FRTLREMRKRSCLGGEFMPSPPTYYKNLKNRVGD
SEQ ID NO:36  QIETYLEHNEGAGVQHLALVTHDI FTTLREMRKRSLGGFEFMPSPPTYYANLHNRAAD
SEQ ID NO:38  QIQTLEHHGGGVQHI AVASSDVLRTLREMRARSAMGGDFLPPPLPKYYEGVRRIAGD
SEQ ID NO:32  QIQTYLEHNEGPGVQHLALVSEDI FNTLREMRKRSGVGGFEFMPSPPLTYKKNLKNRAGD
                                                    301      360

```

FIGURE 2

SEQ ID NO: 41	VLSEAQIKECQELGVLVDRDDQGVLLQIFTKPVGDRPTLFLEMIQRIGCMKDERGEEYQ	
SEQ ID NO: 43	VLSDDDQIKECEEELGILVDRDDQGTLLQIFTKPLGDRPTIFIEIIQRVGCMMKDEEGKAYQ	
SEQ ID NO: 42	VLSDEQIKECEDLGILVDRDDQGTLLQIFTKPVGDRPTLFIEIIQRVGCMLKDDAGQMYQ	
SEQ ID NO: 36	VLTVDDQIKQCEEELGILVDRDDQGTLLQIFTKPVGDRPTIFIEIIQRIGCMVEDEEGKVYQ	
SEQ ID NO: 38	VLSEAQIKECQELGVLVDRDDQGVLLQIFTKPVGDRPTLFLEMIQRIGCMKDERGEEYQ	
SEQ ID NO: 32	VLRDEQIEECEKLGILVDRDDQGTLLQIFTKPVGDRPTLFIEIIQRIGCMKDEQQKLYQ	361 420
SEQ ID NO: 41	KGCGGFGKGNFSELFKSIEDYEKSLKQSAAVQG-S	
SEQ ID NO: 43	SGCGGFGKGNFSELFKSIIEYEKTLKQLVG----	
SEQ ID NO: 42	KGCGGFGKGNFSELFKSIIEYEKTLKQITGSA-AA	
SEQ ID NO: 36	KGACGFGKGNFSELFKSIIEYEKTLKQRAK-----TA.	
SEQ ID NO: 38	KGCGGFGKGNFSELFKSIEDYEKSLKQSAAVQGS.	
SEQ ID NO: 32	KSGCGGFGKGNFSELFKSIIEYEKMLKQVTTETASA.	421 458

Figure 3



SEQUENCE LISTING

<110> Rafalski, J. Antoni
 Cahoon, Rebecca E.
 Coughlan, Sean
 Miao, Guo-Hua

<120> PLANT VITAMIN E BIOSYNTHETIC ENZYMES

<130> BB1289

<140>

<141>

<150> 60/110,781

<151> 1998-12-03

<160> 43

<170> Microsoft Office 97

<210> 1

<211> 792

<212> DNA

<213> Zea mays

<400> 1

```

gttgagccct gttcaagccg agagaggaaa tgctctcgct gcagcgcagg ggttgtcgga 60
tcagggttact ctgcaagttg ctgatgctct ggagcaaccg tttcctgacg ggcagttcga 120
tctgggtgtgg tccatggaga gtggcgagca catgccggac aagagaaaagt ttgttagtga 180
gctagcacgc gtggcggtc ctggaggac aataatcatc gtgacatggt gccataggaa 240
cctggatcca tccgaaacct cgctaaagcc cgatgaactg agcctcctga ggaggatatg 300
cgacgcgtac tacctcccgg actggtgctc accttcagac tatgtgaaca ttgccaagtc 360
actgtctctc gaggatatca agacagctga ctggtcggag aacgtggccc cgttttgcc 420
cgccgtgata aaatcagcgc taacatggaa gggcttcacc tctctgctga cgaccggatg 480
gaagacgatac agaggcgcga tggatgagc gctaataatc cagggctaca agaaggggct 540
catcaaattc accatcatca cctgtcgcaa gcctggagcc gcgtangagg aggcaaggag 600
cacaagttac tagcacagca caggatgcaa gtgcatatgt agatcatggc acatcgccgt 660
cactcatcat actgcacaaa atcaaattctc caggacattt aataattctg cacctcanat 720
attcaggggg gccggtacca atcgccatat gatctatacc gnccacggcg tcnttaactc 780
tnacggaaan ct 792

```

<210> 2

<211> 191

<212> PRT

<213> Zea mays

<400> 2

```

Leu Ser Pro Val Gln Ala Glu Arg Gly Asn Ala Leu Ala Ala Ala Gln
 1              5              10              15

Gly Leu Ser Asp Gln Val Thr Leu Gln Val Ala Asp Ala Leu Glu Gln
 20              25              30

Pro Phe Pro Asp Gly Gln Phe Asp Leu Val Trp Ser Met Glu Ser Gly
 35              40              45

Glu His Met Pro Asp Lys Arg Lys Phe Val Ser Glu Leu Ala Arg Val
 50              55              60

```

Ala Ala Pro Gly Gly Thr Ile Ile Ile Val Thr Trp Cys His Arg Asn
65 70 75 80

Leu Asp Pro Ser Glu Thr Ser Leu Lys Pro Asp Glu Leu Ser Leu Leu
85 90 95

Arg Arg Ile Cys Asp Ala Tyr Tyr Leu Pro Asp Trp Cys Ser Pro Ser
100 105 110

Asp Tyr Val Asn Ile Ala Lys Ser Leu Ser Leu Glu Asp Ile Lys Thr
115 120 125

Ala Asp Trp Ser Glu Asn Val Ala Pro Phe Trp Pro Ala Val Ile Lys
130 135 140

Ser Ala Leu Thr Trp Lys Gly Phe Thr Ser Leu Leu Thr Thr Gly Trp
145 150 155 160

Lys Thr Ile Arg Gly Ala Met Val Met Pro Leu Met Ile Gln Gly Tyr
165 170 175

Lys Lys Gly Leu Ile Lys Phe Thr Ile Ile Thr Cys Arg Lys Pro
180 185 190

<210> 3
<211> 521
<212> DNA
<213> Oryza sativa

<400> 3
cttacagaca aacggcagtt tgtaagcgag ctggcacgag tcgcagctcc tggggcgaga 60
ataatcattg tgacctggtg ccataggaac ctgcagccat ccgaagagtc cctgaaacct 120
gatgagctga atctcctgaa aaggatatgc gatgcatatt atctcccaga ctggtgctct 180
ccttctgatt atgtcaaaat tgccgagtca ctgtctcttg aggatataag gacagctgat 240
tgggtcaagag aacgtcgccc caatccggnc tgcnggttat taaatnaagc aattgacatg 300
gnaagggtta actttctcct ggctaagaan tgggtgggaa gacgattaag aagggtggaat 360
gggtgatgcc tccggatgat nnaaggntac aaagaaangg gtcaacaaat ttaacaanaa 420
caacctgtnc caaagncccg aaacaacgca ataatacccc antaatnaaa ttncgctcct 480
ggctaacctt ctccaacaac gaattaatgg aaantttctga c 521

<210> 4
<211> 82
<212> PRT
<213> Oryza sativa

<400> 4
Leu Thr Asp Lys Arg Gln Phe Val Ser Glu Leu Ala Arg Val Ala Ala
1 5 10 15

Pro Gly Ala Arg Ile Ile Ile Val Thr Trp Cys His Arg Asn Leu Glu
20 25 30

Pro Ser Glu Glu Ser Leu Lys Pro Asp Glu Leu Asn Leu Leu Lys Arg
35 40 45

Ile Cys Asp Ala Tyr Tyr Leu Pro Asp Trp Cys Ser Pro Ser Asp Tyr
50 55 60

Val Lys Ile Ala Glu Ser Leu Ser Leu Glu Asp Ile Arg Thr Ala Asp
65 70 75 80

Trp Ser

<210> 5
 <211> 592
 <212> DNA
 <213> Oryza sativa

<400> 5
 cttacatgta agctcgtgcc gaattcggca cgagcttaca aaatggccca cgccgcccgcg 60
 gccacgggag cactggcacc gctgcatcca ctgctccgct gcacgagccg tcatctctgc 120
 gcctcggttt cccctcgcg cggcctctgc ctccaccacc accgcccgcg ccgcccgcagc 180
 agccggagga cgaaactcgc cgtgcgcgcg atggcaccga cgttgctctc gtcgtcgacg 240
 gcggcggcag ctccccggg gctgaaggag ggcacgcgcg ggctctacga cgaancgtcc 300
 ggcgtgtggg agagcatctg gggcgagcac atgcaccacg gnttctacga cgccggcgag 360
 ggcgccctcca tgtccgacca ccgcccgcgc ccagttcgca tgatcgagga cctcgcccttc 420
 gccgcctccc cgatgatcgg agaagaacca aaatgtattg atttggtgtg gattggtggt 480
 actcaagata ntggngaaca atacggacgc atgctacgna tacttgatcg gtgcagggtga 540
 aagagaaanc ctgcgcgaga caaggtatag caagtcctt taagttgtat ga 592

<210> 6
 <211> 144
 <212> PRT
 <213> Oryza sativa

<400> 6
 Leu Thr Cys Lys Leu Val Pro Asn Ser Ala Arg Ala Tyr Lys Met Ala
 1 5 10 15
 His Ala Ala Ala Thr Gly Ala Leu Ala Pro Leu His Pro Leu Leu
 20 25 30
 Arg Cys Thr Ser Arg His Leu Cys Ala Ser Ala Ser Pro Arg Ala Gly
 35 40 45
 Leu Cys Leu His His His Arg Arg Arg Arg Ser Ser Arg Arg Thr
 50 55 60
 Lys Leu Ala Val Arg Ala Met Ala Pro Thr Leu Ser Ser Ser Ser Thr
 65 70 75 80
 Ala Ala Ala Ala Pro Pro Gly Leu Lys Glu Gly Ile Ala Gly Leu Tyr
 85 90 95
 Asp Glu Xaa Ser Gly Val Trp Glu Ser Ile Trp Gly Glu His Met His
 100 105 110
 His Gly Phe Tyr Asp Ala Gly Glu Gly Ala Ser Met Ser Asp His Arg
 115 120 125
 Arg Ala Pro Val Arg Met Ile Glu Asp Leu Ala Phe Ala Ala Ser Pro
 130 135 140

<210> 7
 <211> 1331
 <212> DNA
 <213> Glycine max

<400> 7

```

gtgacatggc caccgtgggtg aggatcccaa caatctcatg catccacatc cacacgttcc 60
gttcccaatc ccctcgcaact ttcgccagaa tccgggtcgg acccagggtcg tgggctccta 120
ttcggggcatc ggcagcgagc tcggagagag gggagatagt attggagcag aagccgaaga 180
aggatgacaa gaagaagctg cagaaggga tgcgagagtt ttacgacgag tcgtctggct 240
tatgggagaa catttggggc gaccacatgc accatggctt ttatgactcg gattccactg 300
tttcgcttcc ggatcatcgt gctgctcaga tccgaatgat ccaagagtct cttcgctttg 360
cctctgtttc tgaggagcgt agtaaatggc ccaagagtat agttgatgtt ggggtgtggca 420
taggtggcag ctctagatac ctggccaaga aatttggagc aaccagtgtg ggcatactc 480
tgagtcctgt tcaagctcaa agagcaaag ctcttgctgc tgctcaagga ttggctgata 540
aggtttcctt tcaggttgct gacgctctac agcaaccatt ctctgacggc cagtttgatc 600
tgggtgtggc catggagagt ggagagcata tgcctgacaa agctaagttt gttggagagt 660
tagctcgggt agcagcacca ggtgccatta taataatagt aacatgggtc cacagggatc 720
ttggccctga cgaacaatcc ttacatccat gggagcaaga tctcttaaag aagatttgctg 780
atgcatatta cctccctgcc tgggtgctcaa cttctgatta tgttaagttg ctccaatccc 840
tgtcacttta ggacatcaag tcagaagatt ggtctcgctt tgttgctcca ttttgccag 900
cagtgatacg ctcagccttc acatggaag gtctatcttc actcttgagc agtggtaagc 960
ttggaattta tattgcattt caaaaacaaa cccccccatc ttctattgca acttgcaagt 1020
cttatgtcac tgatcattat ttccacacta gataaccctt tacaactaag aacgtagtct 1080
tcatgttcag cgaatatagat aaaaatatgc aacagagtca gagacaggtg gcatgatatt 1140
tacaagaaaa tatcttttat atatataaat gattcaatca aattacttga tgaggattat 1200
gagtgaataa gagaggacag tcatagaaac tttatcctac attccttcta tttccacttc 1260
tgtcaaatat tcctttcatc ttagctatgc tacttgactt gagtaaaaaa aaaaaaaaaa 1320
aaaaaaaaa a 1331

```

<210> 8

<211> 349

<212> PRT

<213> Glycine max

<400> 8

```

Met Ala Thr Val Val Arg Ile Pro Thr Ile Ser Cys Ile His Ile His
  1              5              10             15

Thr Phe Arg Ser Gln Ser Pro Arg Thr Phe Ala Arg Ile Arg Val Gly
          20              25              30

Pro Arg Ser Trp Ala Pro Ile Arg Ala Ser Ala Ala Ser Ser Glu Arg
          35              40              45

Gly Glu Ile Val Leu Glu Gln Lys Pro Lys Lys Asp Asp Lys Lys Lys
  50              55              60

Leu Gln Lys Gly Ile Ala Gly Phe Tyr Asp Glu Ser Ser Gly Leu Trp
  65              70              75              80

Glu Asn Ile Trp Gly Asp His Met His His Gly Phe Tyr Asp Ser Asp
          85              90              95

Ser Thr Val Ser Leu Ser Asp His Arg Ala Ala Gln Ile Arg Met Ile
          100             105             110

Gln Glu Ser Leu Arg Phe Ala Ser Val Ser Glu Glu Arg Ser Lys Trp
          115             120             125

Pro Lys Ser Ile Val Asp Val Gly Cys Gly Ile Gly Gly Ser Ser Arg
          130             135             140

Tyr Leu Ala Lys Lys Phe Gly Ala Thr Ser Val Gly Ile Thr Leu Ser
          145             150             155             160

```

Pro Val Gln Ala Gln Arg Ala Asn Ala Leu Ala Ala Ala Gln Gly Leu
 165 170 175
 Ala Asp Lys Val Ser Phe Gln Val Ala Asp Ala Leu Gln Gln Pro Phe
 180 185 190
 Ser Asp Gly Gln Phe Asp Leu Val Trp Ser Met Glu Ser Gly Glu His
 195 200 205
 Met Pro Asp Lys Ala Lys Phe Val Gly Glu Leu Ala Arg Val Ala Ala
 210 215 220
 Pro Gly Ala Ile Ile Ile Ile Val Thr Trp Cys His Arg Asp Leu Gly
 225 230 235 240
 Pro Asp Glu Gln Ser Leu His Pro Trp Glu Gln Asp Leu Leu Lys Lys
 245 250 255
 Ile Cys Asp Ala Tyr Tyr Leu Pro Ala Trp Cys Ser Thr Ser Asp Tyr
 260 265 270
 Val Lys Leu Leu Gln Ser Leu Ser Leu Gln Asp Ile Lys Ser Glu Asp
 275 280 285
 Trp Ser Arg Phe Val Ala Pro Phe Trp Pro Ala Val Ile Arg Ser Ala
 290 295 300
 Phe Thr Trp Lys Gly Leu Ser Ser Leu Leu Ser Ser Gly Lys Leu Gly
 305 310 315 320
 Ile Tyr Ile Ala Phe Gln Lys Gln Thr Pro Pro Ser Ser Ile Ala Thr
 325 330 335
 Cys Lys Ser Tyr Val Thr Asp His Tyr Phe His Thr Arg
 340 345

<210> 9
 <211> 1011
 <212> DNA
 <213> *Triticum aestivum*

<400> 9
 acgagtcgtc cggcctgtgg gagagcatct ggggcgagca catgcaccac ggcttctacg 60
 actccggcga ggccgcctcc atgtccgacc accgcgcgc ccagatccgc atgatcgagg 120
 aggccctcgc cttcgccgcc gtccccgacg atccgacaaa caaaccacaa acgattgttg 180
 atgttggtatg cggaatcggt ggtagctcaa gatacctggg cgaacaaata tggagcacia 240
 tgctctggga tcacattgac ccagtgcagg ctgagagagg aaatgccctc gcggcagcgc 300
 aaggggttgt cggacaagg ttctttccaa ttgctgatct ctgggagcaa ccatttcctg 360
 gatgggcatt tgatcttgtc cgggnccatg ggagantggt gacacatgcc gaacaaacag 420
 aagtttgtaa gcgagctggc acgcgtcgca gctccaggag caactatcat catcgtgacc 480
 tggtgccata ggaacctcgc gccatcggag gactcactga aacctgacga gctgaatctt 540
 ttgaaaaaga tttgtgatgc atattacctc ccggattggt gctctccctc ggattatgtc 600
 aagattgccg agtcattgtc tcttgaggat atcaaaacgg ccgactggtc tgaaaacgtg 660
 gccccgttct ggctgtgt catccaatca gcactgacat ggaaaggcct cacttctcta 720
 ctaaggagtg gatggaagac gataaaggga gcactgggtga tgcctctcat gatccaaggc 780
 tacaagaaag gcctcattaa gttcaagcat catcacctgc cacaaacccc aagcagccat 840
 agaaggagaa cctggaggcc gcactcgcca agagtggtag aatagaacca tgtgattgga 900
 atagactcng cttgctgtcg ccttggtagc tgaataatc gtgttaccgt gcctctgtat 960
 ctgcaactgg aagtgcata tgagaatggt tcctaaaagc aaaatctcct c 1011

<210> 10
 <211> 293
 <212> PRT
 <213> Triticum aestivum
 .
 <400> 10
 Glu Ser Ser Gly Leu Trp Glu Ser Ile Trp Gly Glu His Met His His
 1 5 10 15
 Gly Phe Tyr Asp Ser Gly Glu Ala Ala Ser Met Ser Asp His Arg Arg
 20 25 30
 Ala Gln Ile Arg Met Ile Glu Glu Ala Leu Ala Phe Ala Ala Val Pro
 35 40 45
 Asp Asp Pro Thr Asn Lys Pro Lys Thr Ile Val Asp Val Gly Cys Gly
 50 55 60
 Ile Gly Gly Ser Ser Arg Tyr Leu Gly Glu Gln Ile Trp Ser Thr Met
 65 70 75 80
 Leu Trp Asp His Ile Asp Pro Val Gln Ala Glu Arg Gly Asn Ala Leu
 85 90 95
 Ala Ala Ala Gln Gly Val Val Arg Thr Arg Phe Phe Pro Ile Ala Asp
 100 105 110
 Leu Trp Glu Gln Pro Phe Pro Gly Trp Ala Phe Asp Leu Val Xaa Xaa
 115 120 125
 Xaa Xaa Xaa Xaa Xaa His Met Pro Asn Lys Gln Lys Phe Val Ser Glu
 130 135 140
 Leu Ala Arg Val Ala Ala Pro Gly Ala Thr Ile Ile Ile Val Thr Trp
 145 150 155 160
 Cys His Arg Asn Leu Ala Pro Ser Glu Asp Ser Leu Lys Pro Asp Glu
 165 170 175
 Leu Asn Leu Leu Lys Lys Ile Cys Asp Ala Tyr Tyr Leu Pro Asp Trp
 180 185 190
 Cys Ser Pro Ser Asp Tyr Val Lys Ile Ala Glu Ser Leu Ser Leu Glu
 195 200 205
 Asp Ile Lys Thr Ala Asp Trp Ser Glu Asn Val Ala Pro Phe Trp Pro
 210 215 220
 Ala Val Ile Gln Ser Ala Leu Thr Trp Lys Gly Leu Thr Ser Leu Leu
 225 230 235 240
 Arg Ser Gly Trp Lys Thr Ile Lys Gly Ala Leu Val Met Pro Leu Met
 245 250 255
 Ile Gln Gly Tyr Lys Lys Gly Leu Ile Lys Phe Lys His His His Leu
 260 265 270
 Pro Gln Thr Pro Ser Ser His Arg Arg Arg Thr Trp Arg Pro His Arg
 275 280 285

Pro Arg Val Val Glu
290

<210> 11
<211> 432
<212> DNA
<213> Oryza sativa

<400> 11
ccacgtcgag ctctggtgag ccgacgccgc gtccgccgcg ggccggttcg ccttcgccct 60
ggcgcgcccg ctgcgcgcca ggteccgacct ctccacgggg aactccgcgc acgectccct 120
cctcctccgc tccgcctccg tcgcgttcct cttcacggcc ccctacggcg gcgaccacgg 180
cgctcgcgcg gacgcggcca ccaccgcctc catcccttcc ttctcccctt cctttctccc 240
cgctcctgga tcaggccaca ggaggggagc gatggtggag gcggccaccg taggcggcgg 300
aggtggcggt cctcctccct agctcccaga cccggctgga ggagggagtg atggtggcgg 360
naggcgngnc tccctcctct cctcctccct tcctcacaat ntggccggag ggaggaangg 420
gccgcggncc aa 432

<210> 12
<211> 75
<212> PRT
<213> Oryza sativa

<400> 12
His Val Glu Leu Trp Cys Ala Asp Ala Ala Ser Ala Ala Gly Arg Phe
1 5 10 15
Ala Phe Ala Leu Gly Ala Pro Leu Ala Ala Arg Ser Asp Leu Ser Thr
20 25 30
Gly Asn Ser Ala His Ala Ser Leu Leu Arg Ser Ala Ser Val Ala
35 40 45
Phe Leu Phe Thr Ala Pro Tyr Gly Gly Asp His Gly Val Gly Ala Asp
50 55 60
Ala Ala Thr Thr Ala Ser Ile Pro Ser Phe Ser
65 70 75

<210> 13
<211> 628
<212> DNA
<213> Oryza sativa

<400> 13
gaagagctac ggcctccgcc ggttcgacca cgtcgtcggc aacgtgccgg agctcgtctc 60
ggtagccgcg tacatctccg ggttcaccgg gttccacgag ttccgccagt tcaccgccga 120
ggacgtgggc accgcccaga gcggcctcaa ctccggtggtg ctccccaaca acgcggagac 180
cgtgctgctg ccgctcaacg agccgggtgca cggcaccaag cgccggagcc agatacagac 240
gtacctggac caccacggcg gcccgggggt gcagcacatc gcgctggcca gcgacgacgt 300
gctcgggacg ctganggaga tgcenggcgc ctccgcatgg gcggttcgat tcttgggccc 360
gccgccgcca actactacga cggtcgcgcg gcgcncggg acttctctcg ggagagcaat 420
taacaatgcc aagactcngg tgtcctggac aaggatacaa gggtttccaa tnttaacaag 480
cattgaanag nnactttctg gngagatcaa gatggtgatg aaagttaatg gaagntncaa 540
aggggntcgc ggttgaaga atntcggctt aatcataggg tngaaacctn agcacagcct 600
anttaggtca gagatgngcn ganaaatt 628

<210> 14
<211> 123

<212> PRT
<213> Oryza sativa

<400> 14
Tyr Gly Leu Arg Arg Phe Asp His Val Val Gly Asn Val Pro Glu Leu
1 5 10 15
Ala Pro Val Ala Ala Tyr Ile Ser Gly Phe Thr Gly Phe His Glu Phe
20 25 30
Ala Glu Phe Thr Ala Glu Asp Val Gly Thr Ala Glu Ser Gly Leu Asn
35 40 45
Ser Val Val Leu Ala Asn Asn Ala Glu Thr Val Leu Leu Pro Leu Asn
50 55 60
Glu Pro Val His Gly Thr Lys Arg Arg Ser Gln Ile Gln Thr Tyr Leu
65 70 75 80
Asp His His Gly Gly Pro Gly Val Gln His Ile Ala Leu Ala Ser Asp
85 90 95
Asp Val Leu Gly Thr Leu Xaa Glu Met Pro Gly Ala Ser Ala Trp Ala
100 105 110
Val Arg Phe Leu Gly Pro Pro Pro Pro Thr Thr
115 120

<210> 15
<211> 1027
<212> DNA
<213> Glycine max

<400> 15
tcacaccaca ccaatgccaa taccatgtg caacgaaatt caagcccaag cccaagccca 60
agcccaagcc caacctgggt ttaagctcgt cggtttcaaa aacttcgtcc gaaccaatcc 120
taagtcggac cgctttcaag tcaaccgctt ccaccacatc gagttctggt gcaccgatgc 180
caccaacgcc tctcgccgat tctcttgagg acttggaatg cctattgttg caaaatctga 240
tctctccacc ggaacccaaa tccacgcctc ctacctcctc cgctccggcg acctctcctt 300
cctcttctcc gctccttact ctccctctct ctcgcccggc tcctccgctg cctcctccgc 360
ctccattccc agtttcgacg ccgccacctg ccttgccctc gctgccaaac acggcttcgg 420
cgctccgcgc atcgcccttg aagtcgccga cgcggaagcc gctttcagcg ccagcgctcg 480
gaaaggagcc gagccggcgt cgcgcgcggt tctcgtcgac gatcgaccg gcttcgcgga 540
ggtgcccctc tacggcgacg tgggtgctccg ctacgtcagc tacaaggacg ccgcgcgcga 600
ggcgccacac gcagatncgt cgcggtggtt cctgcccggg ttcgaggccg cggcgctcgtc 660
gtcttcggtt ccggagctgg actacgggat ccggcggctg gaccacgcg tcgggaacgt 720
tccggagctg gcgcggcggt tgaggtacct gaaaggcttc agcgattcc acgagttcgc 780
ggagttcacc gcggaggacg tgggaacgag cgagagcggg ttgaactcng tggttctggc 840
ngaacaactc ggagacggtg ttgctgccgc tgaacnagcc cggtttacg aacgaaagag 900
gaagaagcca nattgagnnc gtatttngaa cacaanchaa aggtgcttgg tgtgcagcaa 960
ccttgccgnt tgttactcnc naacatcttc ancacactga ggagagatg anaaaanccg 1020
acgtttg 1027

<210> 16
<211> 276
<212> PRT
<213> Glycine max

<400> 16

Met Pro Ile Pro Met Cys Asn Glu Ile Gln Ala Gln Ala Gln Ala Gln
 1 5 10 15
 Ala Gln Ala Gln Pro Gly Phe Lys Leu Val Gly Phe Lys Asn Phe Val
 20 25 30
 Arg Thr Asn Pro Lys Ser Asp Arg Phe Gln Val Asn Arg Phe His His
 35 40 45
 Ile Glu Phe Trp Cys Thr Asp Ala Thr Asn Ala Ser Arg Arg Phe Ser
 50 55 60
 Trp Gly Leu Gly Met Pro Ile Val Ala Lys Ser Asp Leu Ser Thr Gly
 65 70 75 80
 Asn Gln Ile His Ala Ser Tyr Leu Leu Arg Ser Gly Asp Leu Ser Phe
 85 90 95
 Leu Phe Ser Ala Pro Tyr Ser Pro Ser Leu Ser Ala Gly Ser Ser Ala
 100 105 110
 Ala Ser Ser Ala Ser Ile Pro Ser Phe Asp Ala Ala Thr Cys Leu Ala
 115 120 125
 Phe Ala Ala Lys His Gly Phe Gly Val Arg Ala Ile Ala Leu Glu Val
 130 135 140
 Ala Asp Ala Glu Ala Ala Phe Ser Ala Ser Val Ala Lys Gly Ala Glu
 145 150 155 160
 Pro Ala Ser Pro Pro Val Leu Val Asp Asp Arg Thr Gly Phe Ala Glu
 165 170 175
 Val Arg Leu Tyr Gly Asp Val Val Leu Arg Tyr Val Ser Tyr Lys Asp
 180 185 190
 Ala Ala Pro Gln Ala Pro His Ala Asp Xaa Ser Arg Trp Phe Leu Pro
 195 200 205
 Gly Phe Glu Ala Ala Ala Ser Ser Ser Ser Phe Pro Glu Leu Asp Tyr
 210 215 220
 Gly Ile Arg Arg Leu Asp His Ala Val Gly Asn Val Pro Glu Leu Ala
 225 230 235 240
 Pro Ala Val Arg Tyr Leu Lys Gly Phe Ser Gly Phe His Glu Phe Ala
 245 250 255
 Glu Phe Thr Ala Glu Asp Val Gly Thr Ser Glu Ser Gly Leu Asn Ser
 260 265 270
 Val Val Leu Ala
 275

<210> 17

<211> 511

<212> DNA

<213> Vernonia mesipifolia

<400> 17
 ccacaccgat tgccggaact tcaccgcctc tcaccgcctt gcagtcgag caatcgccat 60
 tgaagtcgat gacgccgaat tagctttctc cgtcagcgtc tctcacggcg ctaaaccctc 120
 cgctgctcct gtaacccttg gaaacaacga cgctgtattg tctgaagtta agctttacgg 180
 cgatgtcgct ttccggtaca taagttacaa aaatccgaac tatacatctt cctttttgcc 240
 cgggttcgag cccgttgaaa agacgtcgtc gttttatgac cttgactacg gtatccgccc 300
 tttggaccac gccgtaggaa cgctccctgag cttgcttcgg cagtggacta cgtgaaatca 360
 ttcaccggat tccatgagtt cgccgaattc accgcggagg acgtcgggac gagcgagagg 420
 gaactgaatt cggtcgtttt agcttgcaac agtgagatgg tcttgattcc gatgaacgag 480
 ccggtgtacg gaanaaaagg aagagccaga t 511

<210> 18
 <211> 170
 <212> PRT
 <213> *Vernonia mesipifolia*

<400> 18
 His Thr Asp Cys Arg Asn Phe Thr Ala Ser His Gly Leu Ala Val Arg
 1 5 10 15
 Ala Ile Ala Ile Glu Val Asp Asp Ala Glu Leu Ala Phe Ser Val Ser
 20 25 30
 Val Ser His Gly Ala Lys Pro Ser Ala Ala Pro Val Thr Leu Gly Asn
 35 40 45
 Asn Asp Val Val Leu Ser Glu Val Lys Leu Tyr Gly Asp Val Ala Phe
 50 55 60
 Arg Tyr Ile Ser Tyr Lys Asn Pro Asn Tyr Thr Ser Ser Phe Leu Pro
 65 70 75 80
 Gly Phe Glu Pro Val Glu Lys Thr Ser Ser Phe Tyr Asp Leu Asp Tyr
 85 90 95
 Gly Ile Arg Arg Leu Asp His Ala Val Gly Asn Val Pro Glu Leu Ala
 100 105 110
 Ser Ala Val Asp Tyr Val Lys Ser Phe Thr Gly Phe His Glu Phe Ala
 115 120 125
 Glu Phe Thr Ala Glu Asp Val Gly Thr Ser Glu Arg Glu Leu Asn Ser
 130 135 140
 Val Val Leu Ala Cys Asn Ser Glu Met Val Leu Ile Pro Met Asn Glu
 145 150 155 160
 Pro Val Tyr Gly Xaa Lys Gly Arg Ala Arg
 165 170

<210> 19
 <211> 1165
 <212> DNA
 <213> *Triticum aestivum*

<400> 19
 caagaagcga acacacacca tgccgcccac ccccaccacc cccgcagcca ccggcgccgc 60
 cgcggtgacg ccggagcacg cgcggccgcg ccgaatggtc cgcttcaacc cgcgacgcga 120
 ccgcttcac acgctcgcc tccaccacgt cgagttctgg tgcgcggacg ccgcctccgc 180
 cgccggccgc ttgccttcg cgctcggcgc gccgctcgcc gccaggtccg acctctccac 240

```

ggggaactcc gtgcacgect cccagctgct ccgctcgggc aacctcgect tcctcttcac 300
cgcgccctac gccaacggct gcgacgccgc caccgcctcc ctgccctcct tctccgccga 360
cgccgcgcgc cggttctccg cggaccacgg gctcgcggtg cgctccatag cgctgcgcgt 420
cgcgagacgc gccgagacct tccgcgccag cgtcgacggg ggcgcgcgcc cggccttcag 480
ccccgtggac ctccggccgc gcttcggctt tgcgaggtc gagctctacg gcgacgtcgt 540
gctccgcttc gtcagcatcc ggacggnacg gacgtgcctt cttgccgggg ttcgangggc 600
ttgagcaacc ggggtgccgtg gactaanggc tgacacggnt tgacacgttg tccgnaagtc 660
cggagcttgc ttcggcgccg cctaaccgtg ccggctnaac gggttcaana attcgccagt 720
taacacggag gacgtgggca cggccgagag cgggctcaac tcgatggtgc tcgccaacaa 780
ctcggagggc gtgctgctgc cgctcaacga gccggtgcac ggcaccaagc gccggagcca 840
gatacagacg ttccctggaac accacggcgg ctccggcggtg cagcacatcg cggtggccag 900
cagcgacgtg ctcaggacgc tcaggagat gcgtgcgcgc tccgccatgg gcggcttcga 960
cttctgccca ccccgctgc cgaagtacta cgaaggcgtg cggcgcatcg ccgggatgt 1020
gctctcggag gcgcaaatna aggaatgcaa gaactggggg tgctcntcca caaggagaa 1080
caaagggtgt tgctacaaat cctcaacaag ccaatntggg acaagccgac ttgttctcgt 1140
agatattcac angatctggt gcatg 1165

```

<210> 20

<211> 179

<212> PRT

<213> Triticum aestivum

<400> 20

```

Met Pro Pro Thr Pro Thr Thr Pro Ala Ala Thr Gly Ala Ala Ala Val
  1              5              10              15

Thr Pro Glu His Ala Arg Pro Arg Arg Met Val Arg Phe Asn Pro Arg
      20              25              30

Ser Asp Arg Phe His Thr Leu Ala Phe His His Val Glu Phe Trp Cys
      35              40              45

Ala Asp Ala Ala Ser Ala Ala Gly Arg Phe Ala Phe Ala Leu Gly Ala
      50              55              60

Pro Leu Ala Ala Arg Ser Asp Leu Ser Thr Gly Asn Ser Val His Ala
      65              70              75              80

Ser Gln Leu Leu Arg Ser Gly Asn Leu Ala Phe Leu Phe Thr Ala Pro
      85              90              95

Tyr Ala Asn Gly Cys Asp Ala Ala Thr Ala Ser Leu Pro Ser Phe Ser
      100              105              110

Ala Asp Ala Ala Arg Arg Phe Ser Ala Asp His Gly Leu Ala Val Arg
      115              120              125

Ser Ile Ala Leu Arg Val Ala Asp Ala Ala Glu Ala Phe Arg Ala Ser
      130              135              140

Val Asp Gly Gly Ala Arg Pro Ala Phe Ser Pro Val Asp Leu Gly Arg
      145              150              155              160

Gly Phe Gly Phe Ala Glu Val Glu Leu Tyr Gly Asp Val Val Leu Arg
      165              170              175

Phe Val Ser

```

<210> 21
 <211> 1102
 <212> DNA
 <213> Zea mays

<220>
 <221> unsure
 <222> (454)

<220>
 <221> unsure
 <222> (1072)

<220>
 <221> unsure
 <222> (1083)

<220>
 <221> unsure
 <222> (1092)

<220>
 <221> unsure
 <222> (1100)

<400> 21
 atggctcacg cggcgctgct ccattgctcc cagtcctcca ggagcctcgc agcctgccgc 60
 cgcggcagyc actaccgcgc cccttcgcac gtcccgcgcc actcccgccg tctccgacgc 120
 gccgtcgtca gcctgcgtcc gatggcctcg tcgacggctc agggccccgc gacggcgccg 180
 ccgggtctga aggagggcat cgcggggctg tacgacgagt cgtcggggct gtgggagAAC 240
 atctggggcg accacatgca ccacggcttc tacgactcga gcgaggccgc ctccatggcc 300
 gatcaccgcc gcgcccagat ccgcatgata gaggaggcgc tcgccttcgc cgggtgccca 360
 gcctcagatg atccagagaa gacaccaaaa acaatagtcg atgtcggatg tggcattggt 420
 ggtagctcaa ggtacttggc gaagaaatac ggancgcagt gcactgggat cacgttgagc 480
 cctgttcaag ccgagagagg aaatgctctc gctgcagcgc aggggttgtc ggatcagggt 540
 actctgcaag ttgctgatgc tctggagcaa ccgtttcctg acgggcagtt cgaatctggtg 600
 tgggtccatg agagtggcga gcacatgccg gacaagagaa agtttggttag tgagctagca 660
 cgcgtggcgg ctcttgagg gacaataatc atcgtgacat ggtgccatag gaacctggat 720
 ccacccgaaa cctcgctaaa gcccgatgaa ctgagcctcc tgaggaggat atgcgacgcg 780
 tactacctcc cggactgggtg ctacacctca gactatgtga acattgccaa gtcactgtct 840
 ctcgaggata tcaagacagc tgactggtcg gagaacgtgg ccccgttttg gcccgccgtg 900
 ataaaatcag cgctaactat gaagggttc acctctctgc tgacgaccgg atggaagacg 960
 atcagaggcg cgatggtgat gccgctaata atccagggtc acaagaaggg gctcatcaaa 1020
 ttcaccatca tcacctgtcg caagcctgga gccgcgtagt gatctatacc gnccacggcg 1080
 tcnttaactc tnacggaaan ct 1102

<210> 22
 <211> 352
 <212> PRT
 <213> Zea mays

<220>
 <221> UNSURE
 <222> (152)

<400> 22
 Met Ala His Ala Ala Leu Leu His Cys Ser Gln Ser Ser Arg Ser Leu
 1 5 10 15

Ala Ala Cys Arg Arg Gly Ser His Tyr Arg Ala Pro Ser His Val Pro
 20 25 30
 Arg His Ser Arg Arg Leu Arg Arg Ala Val Val Ser Leu Arg Pro Met
 35 40 45
 Ala Ser Ser Thr Ala Gln Ala Pro Ala Thr Ala Pro Pro Gly Leu Lys
 50 55 60
 Glu Gly Ile Ala Gly Leu Tyr Asp Glu Ser Ser Gly Leu Trp Glu Asn
 65 70 75 80
 Ile Trp Gly Asp His Met His His Gly Phe Tyr Asp Ser Ser Glu Ala
 85 90 95
 Ala Ser Met Ala Asp His Arg Arg Ala Gln Ile Arg Met Ile Glu Glu
 100 105 110
 Ala Leu Ala Phe Ala Gly Val Pro Ala Ser Asp Asp Pro Glu Lys Thr
 115 120 125
 Pro Lys Thr Ile Val Asp Val Gly Cys Gly Ile Gly Gly Ser Ser Arg
 130 135 140
 Tyr Leu Ala Lys Lys Tyr Gly Xaa Gln Cys Thr Gly Ile Thr Leu Ser
 145 150 155 160
 Pro Val Gln Ala Glu Arg Gly Asn Ala Leu Ala Ala Ala Gln Gly Leu
 165 170 175
 Ser Asp Gln Val Thr Leu Gln Val Ala Asp Ala Leu Glu Gln Pro Phe
 180 185 190
 Pro Asp Gly Gln Phe Asp Leu Val Trp Ser Met Glu Ser Gly Glu His
 195 200 205
 Met Pro Asp Lys Arg Lys Phe Val Ser Glu Leu Ala Arg Val Ala Ala
 210 215 220
 Pro Gly Gly Thr Ile Ile Ile Val Thr Trp Cys His Arg Asn Leu Asp
 225 230 235 240
 Pro Ser Glu Thr Ser Leu Lys Pro Asp Glu Leu Ser Leu Leu Arg Arg
 245 250 255
 Ile Cys Asp Ala Tyr Tyr Leu Pro Asp Trp Cys Ser Pro Ser Asp Tyr
 260 265 270
 Val Asn Ile Ala Lys Ser Leu Ser Leu Glu Asp Ile Lys Thr Ala Asp
 275 280 285
 Trp Ser Glu Asn Val Ala Pro Phe Trp Pro Ala Val Ile Lys Ser Ala
 290 295 300
 Leu Thr Trp Lys Gly Phe Thr Ser Leu Leu Thr Thr Gly Trp Lys Thr
 305 310 315 320
 Ile Arg Gly Ala Met Val Met Pro Leu Met Ile Gln Gly Tyr Lys Lys
 325 330 335

Gly Leu Ile Lys Phe Thr Ile Ile Thr Cys Arg Lys Pro Gly Ala Ala
340 345 350

<210> 23
<211> 521
<212> DNA
<213> Oryza sativa

<220>
<221> unsure
<222> (269)

<220>
<221> unsure
<222> (274)

<220>
<221> unsure
<222> (286)

<220>
<221> unsure
<222> (302)

<220>
<221> unsure
<222> (330)

<220>
<221> unsure
<222> (381)..(382)

<220>
<221> unsure
<222> (387)

<220>
<221> unsure
<222> (398)

<220>
<221> unsure
<222> (418)

<220>
<221> unsure
<222> (429)

<220>
<221> unsure
<222> (436)

<220>
<221> unsure
<222> (462)

<220>
<221> unsure
<222> (467)

<220>
 <221> unsure
 <222> (473)

<220>
 <221> unsure
 <222> (514)

<400> 23
 cttacagaca aacggcagtt tgtaagcgag ctggcacgcg tcgcagctcc tggggcgaga 60
 ataatcattg tgacctgggtg ccataggaac ctcgagccat ccgaagagtc cctgaaacct 120
 gatgagctga atctcctgaa aaggatatgc gatgcatatt atctcccaga ctgggtgctct 180
 ccttctgatt atgtcaaaat tgccgagtca ctgtctcttg aggatataag gacagctgat 240
 tggtaagag aacgtcgccc caatccggnc tgcnggttat taaatnaagc aattgacatg 300
 gnaaggggta actttctcct ggctaagaan tgggtgggaa gacgattaag aaggtggaat 360
 gggatgatgcc tccggatgat nnaaggnac aaagaaangg gtcaacaaat ttaacaanaa 420
 caacctgtnc caaagncccg aaacaacgca ataatacccc antaatnaaa ttncgtcct 480
 ggctaacctt ctccaacaac gaattaatgg aaanttctga c 521

<210> 24
 <211> 172
 <212> PRT
 <213> Oryza sativa

<400> 24
 Phe Arg His Gly His Ala Leu Ala Gln Pro Phe Pro Asp Gly Gln Phe
 1 5 10 15
 Asp Leu Val Trp Ser Met Glu Ser Asp Glu His Met Pro Asp Lys Arg
 20 25 30
 Gln Phe Val Ser Glu Leu Ala Arg Val Ala Ala Pro Gly Ala Arg Ile
 35 40 45
 Ile Ile Val Thr Trp Cys His Arg Asn Leu Glu Pro Ser Glu Glu Ser
 50 55 60
 Leu Lys Pro Asp Glu Leu Asn Leu Leu Lys Arg Ile Cys Asp Ala Tyr
 65 70 75 80
 Tyr Leu Pro Asp Trp Cys Ser Pro Ser Asp Tyr Val Lys Ile Ala Glu
 85 90 95
 Ser Leu Ser Leu Glu Asp Ile Arg Thr Ala Asp Trp Ser Glu Asn Val
 100 105 110
 Ala Pro Phe Trp Pro Ala Val Ile Lys Ser Ala Leu Thr Trp Lys Gly
 115 120 125
 Leu Thr Ser Leu Leu Arg Ser Gly Trp Glu Thr Val Arg Gly Ala Met
 130 135 140
 Val Met Pro Leu Val Ile Glu Gly Tyr Lys Lys Gly Leu Ile Lys Phe
 145 150 155 160
 Pro Ile Ile Thr Cys Arg Lys Pro Glu Thr Thr Gln
 165 170

<210> 25
 <211> 464

<212> DNA
<213> Oryza sativa

<400> 25
gcacgagtac agccacggg cgcactggca ccgctgcac cactgctccg ctgcacgagc 60
cgtcatctct gcgcctcggc ttcctctcgc gccggcctct gcctccacca ccaccgccgc 120
cgcccgccga gcagccggag gacgaaactc gccgtgcgcg cgatggcacc gacgttggtc 180
tcgtcgtcga cggcggcggc agctcccccg gggctgaagg agggcatcgc ggggctctac 240
gacgagtcgt ccggcgtgtg ggagagcacc tggggcgagc acatgcacca cgggttctac 300
gacgccggcg aggccgcctc catgtccgac caccgccgcg cccagatccg catgatcgag 360
gaatccctcg ccttcgcccgc cgttccccga tgatgcgggt aacaaaccca aaagtgttat 420
ttactgtttg gtgttgcaaa tgggggtacc tccaaaaaac ttg 464

<210> 26
<211> 128
<212> PRT
<213> Oryza sativa

<400> 26
Ala Arg Val Gln Pro Thr Gly Ala Leu Ala Pro Leu His Pro Leu Leu
1 5 10 15
Arg Cys Thr Ser Arg His Leu Cys Ala Ser Ala Ser Pro Arg Ala Gly
20 25 30
Leu Cys Leu His His His Arg Arg Arg Arg Ser Ser Arg Arg Thr
35 40 45
Lys Leu Ala Val Arg Ala Met Ala Pro Thr Leu Ser Ser Ser Ser Thr
50 55 60
Ala Ala Ala Ala Pro Pro Gly Leu Lys Glu Gly Ile Ala Gly Leu Tyr
65 70 75 80
Asp Glu Ser Ser Gly Val Trp Glu Ser Ile Trp Gly Glu His Met His
85 90 95
His Gly Phe Tyr Asp Ala Gly Glu Ala Ala Ser Met Ser Asp His Arg
100 105 110
Arg Ala Gln Ile Arg Met Ile Glu Glu Ser Leu Ala Phe Ala Ala Val
115 120 125

<210> 27
<211> 1189
<212> DNA
<213> Glycine max

<400> 27
ggacatggcc accgtggtga ggatcccaac aatctcatgc atccacatcc acacgttccg 60
ttcccaatcc cctcgcactt tcgccagaat ccgggtcggg cccaggtcgt gggctcctat 120
tcgggcatcg gcagcgagct cggagagagg ggagatagta ttggagcaga agccgaagaa 180
ggatgacaag aagaagctgc agaagggaat cgcagagttt tacgacgagt cgtctggctt 240
atgggagaac atttggggcg accacatgca ccatggcttt tatgactcgg attccactgt 300
ttcgttttcg gatcatcgtg ctgctcagat ccgaatgatc caagagtctc ttcgctttgc 360
ctctgtttct gaggagcgta gtaaattggc caagagtata gttgatgttg ggtgtggcat 420
aggtggcagc tctagatacc tggccaagaa atttgagca accagtgtag gcactactct 480
gagtcctggt caagctcaaa gagcaaatgc tcttgctgct gctcaaggat tggctgataa 540
ggtttctctt caggttgctg acgctctaca gcaaccattc tctgacggcc agtttgatct 600
ggtgtggtcc atggagagtg gagagcatat gcctgacaaa gctaagtttg ttggagagtt 660

```

agctcgggta gcagcaccag gtgccattat aataatagta acatgggtgcc acagggatct 720
tggccctgac gaacaatcct tacatccatg ggagcaagat ctcttaaaga agatttgcca 780
tgcattattac ctccctgcct ggtgctcaac ttctgattat gtttaagttgc tccaatccct 840
gtcactttcag gacatcaagt cagaagattg gtctcgcttt gttgctccat tttggccagc 900
agtgtatcgc tcagccttca catggaagggt tctatcttca ctcttgagca gtggacaaaa 960
aacgataaaa ggagctttgg ctatgccatt gatgatagag ggatacaaga aagatctaata 1020
taagtttgcc atcattacat gtcgaaaacc tgaataaatg gagaggcagg attactttta 1080
tagaatgaac caagtttcca acaggtcggt tatttcgata gttgagaaac aagagaaaaa 1140
ataaatgaaa ggggttggtc gattttaaaa aaaaaaaaaa aaaaaaaaaa 1189

```

<210> 28
 <211> 350
 <212> PRT
 <213> Glycine max

<400> 28
 Met Ala Thr Val Val Arg Ile Pro Thr Ile Ser Cys Ile His Ile His
 1 5 10 15
 Thr Phe Arg Ser Gln Ser Pro Arg Thr Phe Ala Arg Ile Arg Val Gly
 20 25 30
 Pro Arg Ser Trp Ala Pro Ile Arg Ala Ser Ala Ala Ser Ser Glu Arg
 35 40 45
 Gly Glu Ile Val Leu Glu Gln Lys Pro Lys Lys Asp Asp Lys Lys Lys
 50 55 60
 Leu Gln Lys Gly Ile Ala Glu Phe Tyr Asp Glu Ser Ser Gly Leu Trp
 65 70 75 80
 Glu Asn Ile Trp Gly Asp His Met His His Gly Phe Tyr Asp Ser Asp
 85 90 95
 Ser Thr Val Ser Leu Ser Asp His Arg Ala Ala Gln Ile Arg Met Ile
 100 105 110
 Gln Glu Ser Leu Arg Phe Ala Ser Val Ser Glu Glu Arg Ser Lys Trp
 115 120 125
 Pro Lys Ser Ile Val Asp Val Gly Cys Gly Ile Gly Gly Ser Ser Arg
 130 135 140
 Tyr Leu Ala Lys Lys Phe Gly Ala Thr Ser Val Gly Ile Thr Leu Ser
 145 150 155 160
 Pro Val Gln Ala Gln Arg Ala Asn Ala Leu Ala Ala Ala Gln Gly Leu
 165 170 175
 Ala Asp Lys Val Ser Phe Gln Val Ala Asp Ala Leu Gln Gln Pro Phe
 180 185 190
 Ser Asp Gly Gln Phe Asp Leu Val Trp Ser Met Glu Ser Gly Glu His
 195 200 205
 Met Pro Asp Lys Ala Lys Phe Val Gly Glu Leu Ala Arg Val Ala Ala
 210 215 220
 Pro Gly Ala Ile Ile Ile Ile Val Thr Trp Cys His Arg Asp Leu Gly
 225 230 235 240

$\langle 210 \rangle$	30
$\langle 211 \rangle$	366

<212> PRT
 <213> Triticum aestivum

<220>
 <221> UNSURE
 <222> (5)

<220>
 <221> UNSURE
 <222> (45)

<400> 30

```

Met Ala Asn Ser Xaa Arg Pro Ala Pro Leu Thr Pro Leu His Arg Leu
 1              5              10              15

Asp Ala Ala Pro Pro Pro Arg Pro Ser Leu Gly His Ala Ala Arg Pro
      20              25              30

Val Pro Arg Pro Val Leu Pro Leu Leu Pro Ala Arg Xaa Leu Arg Ala
      35              40              45

Pro Asp Gly Val Val Asp Asp Arg Gly Pro Gly Asp Ala Ala Pro Pro
      50              55              60

Gly Leu Lys Glu Gly Ile Ala Gly Leu Tyr Asp Glu Ser Ser Gly Leu
 65              70              75              80

Trp Glu Ser Ile Trp Gly Glu His Met His His Gly Phe Tyr Asp Ser
      85              90              95

Gly Glu Ala Ala Ser Met Ser Asp His Arg Arg Ala Gln Ile Arg Met
      100              105              110

Ile Glu Glu Ala Leu Ala Phe Ala Ala Val Pro Asp Asp Pro Thr Asn
      115              120              125

Lys Pro Lys Thr Ile Val Asp Val Gly Cys Gly Ile Gly Gly Ser Ser
      130              135              140

Arg Tyr Leu Ala Asn Lys Tyr Gly Ala Gln Cys Ser Gly Ile Thr Leu
      145              150              155              160

Ser Pro Val Gln Ala Glu Arg Gly Asn Ala Leu Ala Ala Ala Gln Gly
      165              170              175

Leu Ser Asp Lys Ala Ser Phe Gln Val Ala Asp Ala Leu Glu Gln Pro
      180              185              190

Phe Pro Asp Gly Gln Phe Asp Leu Val Trp Ser Met Glu Ser Gly Glu
      195              200              205

His Met Pro Asn Lys Gln Lys Phe Val Ser Glu Leu Ala Arg Val Ala
      210              215              220

Ala Pro Gly Ala Thr Ile Ile Ile Val Thr Trp Cys His Arg Asn Leu
      225              230              235              240

Ala Pro Ser Glu Asp Ser Leu Lys Pro Asp Glu Leu Asn Leu Leu Lys
      245              250              255
  
```

Lys Ile Cys Asp Ala Tyr Tyr Leu Pro Asp Trp Cys Ser Pro Ser Asp
260 265 270

Tyr Val Lys Ile Ala Glu Ser Leu Ser Leu Glu Asp Ile Lys Thr Ala
275 280 285

Asp Trp Ser Glu Asn Val Ala Pro Phe Trp Pro Ala Val Ile Gln Ser
290 295 300

Ala Leu Thr Trp Lys Gly Leu Thr Ser Leu Leu Arg Ser Gly Trp Lys
305 310 315 320

Thr Ile Lys Gly Ala Leu Val Met Pro Leu Met Ile Gln Gly Tyr Lys
325 330 335

Lys Gly Leu Ile Lys Phe Ser Ile Ile Thr Cys Arg Lys Pro Gln Ala
340 345 350

Ala Ile Glu Gly Glu Pro Glu Ala Ala Ser Pro Ser Val Glu
355 360 365

<210> 31
<211> 1605
<212> DNA
<213> Catalpa sp.

<400> 31
gcacgaggca ccacttcacc accaccacca ccaccaccac caccacaact gcttccaccg 60
ccgcagagtc actctatcag ttaagatcaa agcaacaagt tcagaatccc aaacaatggg 120
caagcagacg accacttccg ccaccgccgc ggacgggtcc aaagatgcgc atgcagaatt 180
caagctgggtg ggcttcaaga atttcgtcag gaccaacccc aagtccgacc acttctgcgt 240
ccaccgcttc caccatatag agttctggtg cggcgacgcc accaacaccg ccaagcgctt 300
ctcttggggc ctcggtatgc ccctcgtcgc caaatcggat cttccactg gaaactccgc 360
tcatgcctcg tatcttcttc ggtctggcga actcaacttc ctcttcacga gcccttactc 420
tccttcaatc tccgcccct cctccgccgc catcccagtt ttctccttct ccacctacca 480
gtcttttacc tctcccatg gcctcgctgt tcgtgcggtg gctattcagg tcgattcggc 540
cttttcggct tactctgcct ccatttcccg cggcgccaaa cccgtgtccg caccgattct 600
tttatctgac aacaagactg ccattgcgga ggttcattta tatggagact cagtgttgcg 660
attcgtgagc tatggtgata atgggacag cccagatgga tggttcttgc cgggctttga 720
gcctgtggat gatcagatgt cttataaaga attggattat gggattagaa ggctagatca 780
tgctgtagga aatgtgcccg aactcgggtc agttgtggat tacttgaaaa aattcacagg 840
gtttcatgaa tttgcagagt ttacttcaga ggatgtggga acagcagaaa gtggattgaa 900
ttctatgggt ttagcgaaca acaatgaaaa tgtgttgta cctctgaacg aaccgggtgt 960
tggaaccaag aggaagagcc agattcagac ttatttggag cacaatgaag ggccagggtg 1020
acagcatttg gcattagtga gtgaggatat ctttaacaca ttaagggaaa tgagaaagag 1080
gagtggagtt gggggattcg agttcatgcc ttgcctccg cttacttatt acaagaatct 1140
caagaacaga gctggagatg tgctgaggga tgagcagatt gaggagtgtg agaagttggg 1200
gatcttggtg gacagggatg atcaggggac tttgcttcag attttcacca agcctgtggg 1260
tgataggcca acgctattca tagagatcat tcagagaatc ggggtgcattc tcaaagacga 1320
acaaggaag ctctaccaga agagtgggtg tggaggattt ggaaagggca acttctccga 1380
actcttcaaa tccatcgaag aatacagaaa aatgctcgaa gcaaagcaag tcaactgaaac 1440
agcgtcggcc tgagttctga gtccttctta ctgtgttgta gatatgttga tgaaccaatg 1500
tcctgtcggg acataggttg ttcttatgct gtactaaact gtagttgaca agaagtttta 1560
cttaataata tatcttactt tctataaaaa aaaaaaaaaa aaaaa 1605

<210> 32
<211> 445
<212> PRT
<213> Catalpa sp.

<400> 32
 Met Gly Lys Gln Thr Thr Thr Ser Ala Thr Ala Ala Asp Gly Ser Lys
 1 5 10 15
 Asp Ala His Ala Glu Phe Lys Leu Val Gly Phe Lys Asn Phe Val Arg
 20 25 30
 Thr Asn Pro Lys Ser Asp His Phe Cys Val His Arg Phe His His Ile
 35 40 45
 Glu Phe Trp Cys Gly Asp Ala Thr Asn Thr Ala Lys Arg Phe Ser Trp
 50 55 60
 Gly Leu Gly Met Pro Leu Val Ala Lys Ser Asp Leu Ser Thr Gly Asn
 65 70 75 80
 Ser Ala His Ala Ser Tyr Leu Leu Arg Ser Gly Glu Leu Asn Phe Leu
 85 90 95
 Phe Thr Ser Pro Tyr Ser Pro Ser Ile Ser Ala Pro Ser Ser Ala Ala
 100 105 110
 Ile Pro Ser Phe Ser Phe Ser Thr Tyr Gln Ser Phe Thr Ser Ser His
 115 120 125
 Gly Leu Ala Val Arg Ala Val Ala Ile Gln Val Asp Ser Ala Phe Ser
 130 135 140
 Ala Tyr Ser Ala Ser Ile Ser Arg Gly Ala Lys Pro Val Ser Ala Pro
 145 150 155 160
 Ile Leu Leu Ser Asp Asn Lys Thr Ala Ile Ala Glu Val His Leu Tyr
 165 170 175
 Gly Asp Ser Val Leu Arg Phe Val Ser Tyr Gly Asp Asn Gly Thr Gly
 180 185 190
 Pro Asp Gly Trp Phe Leu Pro Gly Phe Glu Pro Val Asp Asp Gln Met
 195 200 205
 Ser Tyr Lys Glu Leu Asp Tyr Gly Ile Arg Arg Leu Asp His Ala Val
 210 215 220
 Gly Asn Val Pro Glu Leu Gly Pro Val Val Asp Tyr Leu Lys Lys Phe
 225 230 235 240
 Thr Gly Phe His Glu Phe Ala Glu Phe Thr Ser Glu Asp Val Gly Thr
 245 250 255
 Ala Glu Ser Gly Leu Asn Ser Met Val Leu Ala Asn Asn Asn Glu Asn
 260 265 270
 Val Leu Leu Pro Leu Asn Glu Pro Val Phe Gly Thr Lys Arg Lys Ser
 275 280 285
 Gln Ile Gln Thr Tyr Leu Glu His Asn Glu Gly Pro Gly Val Gln His
 290 295 300
 Leu Ala Leu Val Ser Glu Asp Ile Phe Asn Thr Leu Arg Glu Met Arg
 305 310 315 320

Lys Arg Ser Gly Val Gly Gly Phe Glu Phe Met Pro Ser Pro Pro Leu
 325 330 335
 Thr Tyr Tyr Lys Asn Leu Lys Asn Arg Ala Gly Asp Val Leu Arg Asp
 340 345 350
 Glu Gln Ile Glu Glu Cys Glu Lys Leu Gly Ile Leu Val Asp Arg Asp
 355 360 365
 Asp Gln Gly Thr Leu Leu Gln Ile Phe Thr Lys Pro Val Gly Asp Arg
 370 375 380
 Pro Thr Leu Phe Ile Glu Ile Ile Gln Arg Ile Gly Cys Met Leu Lys
 385 390 395 400
 Asp Glu Gln Gly Lys Leu Tyr Gln Lys Ser Gly Cys Gly Gly Phe Gly
 405 410 415
 Lys Gly Asn Phe Ser Glu Leu Phe Lys Ser Ile Glu Glu Tyr Glu Lys
 420 425 430
 Met Leu Glu Ala Lys Gln Val Thr Glu Thr Ala Ser Ala
 435 440 445

<210> 33
 <211> 1106
 <212> DNA
 <213> Oryza sativa

<400> 33
 gcacgaggaa gagctacggc ctccgccggt tcgaccacgt cgtcggcaac gtgccggagc 60
 tcgctccggt agccgcgtac atctccgggt tcaccgggtt ccacgagttc gccgagttca 120
 ccgccgagga cgtggggcacc gccgagagcg gcctcaactc ggtgggtgctc gccacaacg 180
 cggagaccgt gctgctgccg ctcaacgagc cggtgacagg caccaagcgg cggagccaga 240
 tacagacgta cctggaccac caccggcgcc cgggggtgca gcacatcgcg ctggccagcg 300
 acgacgtgct cgggacgctg agggagatgc gggcgcgctc cgccatgggc ggcttcgagt 360
 tcttgggcgcc gccgccgccc aactactacg acggcgtgcg gcggcgcgcc ggggacgtgc 420
 tctcggagga gcagatcaac gagtgccagg agctcggggt gctcgtggac agggatgacc 480
 agggggtggt gctccagatc ttcaccaagc cagtaggaga caggccaacc ttttcttgg 540
 agatgataca aaggattggg tgcattggga aggatgagag tgggcaggag taccagaagg 600
 gcggctgctg cgggtttggg aagggaact tctcggagct gttcaagtcc attgaggagt 660
 atgagaaatc cttgaagcc aagcaagccc ctacagttca aggatcctag gtaggaactg 720
 gaggcctgga gcaacagatg taaccagtgt atttgtatta tggagcagaa gaaaaaagat 780
 gtgctttcac tgctttgtga tatgtgtcat gcaagttgat gttgtaattt gtggaagctg 840
 aagacaaatg atggtacaat cactgtaata gataatagac atggatcaca tacaagaatg 900
 taacctagtg ttggcattgc tgctgtacaa tcttgcttgg aaataaaata ataatacaacc 960
 tggagaaaga atgtaacctg ctgttgcat tgctgatgta caatcttttt ttggaaataa 1020
 aataagaatc cccccaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 1080
 aaaaaaaaaa aaaaaaaaaa aaaaaa 1106

<210> 34
 <211> 235
 <212> PRT
 <213> Oryza sativa

<400> 34
 Thr Arg Lys Ser Tyr Gly Leu Arg Arg Phe Asp His Val Val Gly Asn
 1 5 10 15

Val Pro Glu Leu Ala Pro Val Ala Ala Tyr Ile Ser Gly Phe Thr Gly
 20 25 30
 Phe His Glu Phe Ala Glu Phe Thr Ala Glu Asp Val Gly Thr Ala Glu
 35 40 45
 Ser Gly Leu Asn Ser Val Val Leu Ala Asn Asn Ala Glu Thr Val Leu
 50 55 60
 Leu Pro Leu Asn Glu Pro Val His Gly Thr Lys Arg Arg Ser Gln Ile
 65 70 75 80
 Gln Thr Tyr Leu Asp His His Gly Gly Pro Gly Val Gln His Ile Ala
 85 90 95
 Leu Ala Ser Asp Asp Val Leu Gly Thr Leu Arg Glu Met Arg Ala Arg
 100 105 110
 Ser Ala Met Gly Gly Phe Glu Phe Leu Ala Pro Pro Pro Pro Asn Tyr
 115 120 125
 Tyr Asp Gly Val Arg Arg Arg Ala Gly Asp Val Leu Ser Glu Glu Gln
 130 135 140
 Ile Asn Glu Cys Gln Glu Leu Gly Val Leu Val Asp Arg Asp Asp Gln
 145 150 155 160
 Gly Val Leu Leu Gln Ile Phe Thr Lys Pro Val Gly Asp Arg Pro Thr
 165 170 175
 Phe Phe Leu Glu Met Ile Gln Arg Ile Gly Cys Met Glu Lys Asp Glu
 180 185 190
 Ser Gly Gln Glu Tyr Gln Lys Gly Gly Cys Gly Gly Phe Gly Lys Gly
 195 200 205
 Asn Phe Ser Glu Leu Phe Lys Ser Ile Glu Glu Tyr Glu Lys Ser Leu
 210 215 220
 Glu Ala Lys Gln Ala Pro Thr Val Gln Gly Ser
 225 230 235

<210> 35
 <211> 1550
 <212> DNA
 <213> Glycine max

<400> 35
 tcacaccaca ccaatgccaa taccatgtg caacgaaatt caagcccaag cccaagccca 60
 agcccaagcc caacctgggt ttaagctcgt cggtttcaaa aacttcgtcc gaaccaatcc 120
 taagtcggac cgctttcaag tcaaccgctt ccaccacatc gagttctggt gcaccgatgc 180
 caccaacgcc tctcgccgat tctcttgagg acttggaatg cctattgtgg caaaatctga 240
 tctctccacc ggaaaccaa tccacgcctc ctacctctc cgctccggcg acctctcctt 300
 cctcttctcc gctccttact ctccctctct ctccgcccgc tctccgctg cctcctccgc 360
 ctccattccc agtttcgacg ccgccacctg ccttgccctc gctgccaaac acggttcgg 420
 cgcccgccgc atcgcccttg aagtcgccga cgcggaagcc gctttcagcg ccagcgtcgc 480
 gaaaggagcc gagccggcgt cgcgcgcggt tctcgtcgac gatcgaccg gcttcgcgga 540
 ggtgcgcctc tacggcgacg tgggtctccg ctacgtcagc tacaaggacg ccgcgcgcga 600
 ggccgacac gcagatccgt cgcggtggtt cctgccggga ttcgaggccg cggcgtcgtc 660
 gtcttcggtt ccggagctgg actacgggat ccggcggtg gaccacgccg tcgggaacgt 720


```

tccggagctg ggcgcggcgg tgaggtacct gaaaggcttc agcggattcc acgagttcgc 780
ggagttcacc gcgaggagcg tgggaacgag cgagagcggg ttgaactcgg tggttctggc 840
gaacaactcg gagacggtgt tgctgccgct gaacgagccg gtttacggaa cgaagaggaa 900
gagccagatt gagacgtatt tggaaacaaa cgaagggtgt ggtgtgcagc accttgcgct 960
tgttactcac gacatcttca ccacactgag agagatgaga aagcgaagtt tccttggtgg 1020
at ttgagttc atgccttctc ctccctccac ctattacgcc aacctccaca accgtgccgc 1080
tgatgtgttg accgttgacc agattaagca gtgtgaggag cttgggattc ttgttgacag 1140
agatgatcag ggcactctgc ttcagatttt caccaagcct gttggggaca ggccaacgat 1200
attcatagag ataattcaga ggcacgggtg catggtggag gatgaggaag ggaagggtga 1260
ccagaagggt gcatgtgggg gttttgggaa aggcattttt tctgagcttt tcaaattccat 1320
tgaagaatat gagaagactt tggaagctaa aagaaccgag taagcacatt ggaagaacac 1380
aaatactcct ttgttgaaat gattaatgag gaatcaatgt ggcatagggt gtttatactc 1440
tataatacat agaattacaa tgatagtgtc ctcccttgta tgaaaaatgaa atcacagaaa 1500
cttttatgga tagtattttt ctattaaaaa aaaaaaaaaa aaaaaaaaaa 1550

```

<210> 36
 <211> 449
 <212> PRT
 <213> Glycine max

<400> 36
 Met Pro Ile Pro Met Cys Asn Glu Ile Gln Ala Gln Ala Gln Ala Gln
 1 5 10 15
 Ala Gln Ala Gln Pro Gly Phe Lys Leu Val Gly Phe Lys Asn Phe Val
 20 25 30
 Arg Thr Asn Pro Lys Ser Asp Arg Phe Gln Val Asn Arg Phe His His
 35 40 45
 Ile Glu Phe Trp Cys Thr Asp Ala Thr Asn Ala Ser Arg Arg Phe Ser
 50 55 60
 Trp Gly Leu Gly Met Pro Ile Val Ala Lys Ser Asp Leu Ser Thr Gly
 65 70 75 80
 Asn Gln Ile His Ala Ser Tyr Leu Leu Arg Ser Gly Asp Leu Ser Phe
 85 90 95
 Leu Phe Ser Ala Pro Tyr Ser Pro Ser Leu Ser Ala Gly Ser Ser Ala
 100 105 110
 Ala Ser Ser Ala Ser Ile Pro Ser Phe Asp Ala Ala Thr Cys Leu Ala
 115 120 125
 Phe Ala Ala Lys His Gly Phe Gly Val Arg Ala Ile Ala Leu Glu Val
 130 135 140
 Ala Asp Ala Glu Ala Ala Phe Ser Ala Ser Val Ala Lys Gly Ala Glu
 145 150 155 160
 Pro Ala Ser Pro Pro Val Leu Val Asp Asp Arg Thr Gly Phe Ala Glu
 165 170 175
 Val Arg Leu Tyr Gly Asp Val Val Leu Arg Tyr Val Ser Tyr Lys Asp
 180 185 190
 Ala Ala Pro Gln Ala Pro His Ala Asp Pro Ser Arg Trp Phe Leu Pro
 195 200 205

Gly Phe Glu Ala Ala Ala Ser Ser Ser Ser Phe Pro Glu Leu Asp Tyr
 210 215 220
 Gly Ile Arg Arg Leu Asp His Ala Val Gly Asn Val Pro Glu Leu Ala
 225 230 235 240
 Pro Ala Val Arg Tyr Leu Lys Gly Phe Ser Gly Phe His Glu Phe Ala
 245 250 255
 Glu Phe Thr Ala Glu Asp Val Gly Thr Ser Glu Ser Gly Leu Asn Ser
 260 265 270
 Val Val Leu Ala Asn Asn Ser Glu Thr Val Leu Leu Pro Leu Asn Glu
 275 280 285
 Pro Val Tyr Gly Thr Lys Arg Lys Ser Gln Ile Glu Thr Tyr Leu Glu
 290 295 300
 His Asn Glu Gly Ala Gly Val Gln His Leu Ala Leu Val Thr His Asp
 305 310 315 320
 Ile Phe Thr Thr Leu Arg Glu Met Arg Lys Arg Ser Phe Leu Gly Gly
 325 330 335
 Phe Glu Phe Met Pro Ser Pro Pro Pro Thr Tyr Tyr Ala Asn Leu His
 340 345 350
 Asn Arg Ala Ala Asp Val Leu Thr Val Asp Gln Ile Lys Gln Cys Glu
 355 360 365
 Glu Leu Gly Ile Leu Val Asp Arg Asp Asp Gln Gly Thr Leu Leu Gln
 370 375 380
 Ile Phe Thr Lys Pro Val Gly Asp Arg Pro Thr Ile Phe Ile Glu Ile
 385 390 395 400
 Ile Gln Arg Ile Gly Cys Met Val Glu Asp Glu Glu Gly Lys Val Tyr
 405 410 415
 Gln Lys Gly Ala Cys Gly Gly Phe Gly Lys Gly Asn Phe Ser Glu Leu
 420 425 430
 Phe Lys Ser Ile Glu Glu Tyr Glu Lys Thr Leu Glu Ala Lys Arg Thr
 435 440 445

Ala

<210> 37
 <211> 1614
 <212> DNA
 <213> Triticum aestivum

<400> 37
 gcacgagcaa gaagcgaaca cacaccatgc cgcccccccc caccaccccc gcagccaccg 60
 gcgcccgcgc ggtgacgccg gagcacgcgc ggccgcgcgc aatgggtccgc ttcaaccgcg 120
 gcagcgaccg cttccacacg ctgccttcc accacgtcga gttctggtgc gcggacgcgc 180
 cctccgcgcgc cgccgcgttc gccttcgcgc tcggcgcgcc gctcgccgcc aggtccgacc 240
 tctccacggg gaactccgtg cagcctccc agctgctccg ctggggcaac ctgccttcc 300
 tcttcacggc cccctacgcc aacggctgcg acgcccgcac cgctccctg ccctccttct 360

```

ccgccgacgc cgcgcgccag ttctccgcgc accacggcct cgcgggtgcgc tccatagcgc 420
tgcgcgtgcgc ggacgctgcc gaggccttcc gcgccagcgt cgacgggggc gcgcgcccg 480
ccttcagccc tgtggacctc ggccgcggct tcggcttcgc ggaggtcgag ctctacggcg 540
acgtcgtgct ccgcttcgct agccaccggc acggcaggga cgtgcccttc ttgccggggt 600
tcgagggcgt gagcaacca gacgccgtgg actacggcct gacgcgggtc gaccacgtcg 660
tcggcaacgt cccggagctt gccccgcgc cgccctacgt cgcggggttc acggggttc 720
acgagttcgc cgagttcacg acggaggacg tgggcacggc cgagagcggg ctcaactcga 780
tgggtgctgc caacaactcg gagggcgtgc tgctgccgt caacgagccg gtgcacggca 840
ccaagcgccg gagccagata cagacgttcc tggaaacca cggcggtcg ggcgtgcagc 900
acatcgccgt ggccagcagc gacgtgctca ggacgtcag ggagatgctg gcgcgtccg 960
ccatgggcgg cttcgacttc ctgccacccc cgctgccgaa gtactacgaa ggcgtgcggc 1020
gcatcgccgg ggtgtgctc tcggagcgcc agatcaagga atgccaggag ctgggggtgc 1080
tcgtcgacag ggacgaccaa ggggtgtgct tacaatctt caccaagcca gtaggggaca 1140
ggccgacgtt gttcctggag atgatccaga ggtcgggtg catggagaag gacgagagag 1200
gggaagagta ccagaagggt ggctgcggcg gggtcggcaa aggcacttc tccgagctgt 1260
tcaagtccat tgaagattac gagaagtccc ttgaagccaa gcaatctgct gcagttcagg 1320
gatcatagga tagaagctgg agctggaggà gctgatccag tactttgtat caggtctcat 1380
ggagcaaaag aaaatgatgt tgtttgtaag atgcggcgcg caattatgtc cgatgttata 1440
attgtgaag ctgaagacag atgtatccta tgtatgatgg gtgtaataga tggtagaggg 1500
ggctcggtc acacatgaac aaaatgtact gttggcattg ttgtataatc ttgcttgcaa 1560
gtaaaataaa gaagaaccga ttttgagttc tgcataaaaa aaaaaaaaaa aaaa 1614

```

<210> 38
 <211> 433
 <212> PRT
 <213> Triticum aestivum

<400> 38
 Met Pro Pro Thr Pro Thr Thr Pro Ala Ala Thr Gly Ala Ala Ala Val
 1 5 10 15
 Thr Pro Glu His Ala Arg Pro Arg Arg Met Val Arg Phe Asn Pro Arg
 20 25 30
 Ser Asp Arg Phe His Thr Leu Ala Phe His His Val Glu Phe Trp Cys
 35 40 45
 Ala Asp Ala Ala Ser Ala Ala Gly Arg Phe Ala Phe Ala Leu Gly Ala
 50 55 60
 Pro Leu Ala Ala Arg Ser Asp Leu Ser Thr Gly Asn Ser Val His Ala
 65 70 75 80
 Ser Gln Leu Leu Arg Ser Gly Asn Leu Ala Phe Leu Phe Thr Ala Pro
 85 90 95
 Tyr Ala Asn Gly Cys Asp Ala Ala Thr Ala Ser Leu Pro Ser Phe Ser
 100 105 110
 Ala Asp Ala Ala Arg Gln Phe Ser Ala Asp His Gly Leu Ala Val Arg
 115 120 125
 Ser Ile Ala Leu Arg Val Ala Asp Ala Ala Glu Ala Phe Arg Ala Ser
 130 135 140
 Val Asp Gly Gly Ala Arg Pro Ala Phe Ser Pro Val Asp Leu Gly Arg
 145 150 155 160
 Gly Phe Gly Phe Ala Glu Val Glu Leu Tyr Gly Asp Val Val Leu Arg
 165 170 175

Phe Val Ser His Pro Asp Gly Arg Asp Val Pro Phe Leu Pro Gly Phe
 180 185 190
 Glu Gly Val Ser Asn Pro Asp Ala Val Asp Tyr Gly Leu Thr Arg Phe
 195 200 205
 Asp His Val Val Gly Asn Val Pro Glu Leu Ala Pro Ala Ala Ala Tyr
 210 215 220
 Val Ala Gly Phe Thr Gly Phe His Glu Phe Ala Glu Phe Thr Thr Glu
 225 230 235 240
 Asp Val Gly Thr Ala Glu Ser Gly Leu Asn Ser Met Val Leu Ala Asn
 245 250 255
 Asn Ser Glu Gly Val Leu Leu Pro Leu Asn Glu Pro Val His Gly Thr
 260 265 270
 Lys Arg Arg Ser Gln Ile Gln Thr Phe Leu Glu His His Gly Gly Ser
 275 280 285
 Gly Val Gln His Ile Ala Val Ala Ser Ser Asp Val Leu Arg Thr Leu
 290 295 300
 Arg Glu Met Arg Ala Arg Ser Ala Met Gly Gly Phe Asp Phe Leu Pro
 305 310 315 320
 Pro Pro Leu Pro Lys Tyr Tyr Glu Gly Val Arg Arg Ile Ala Gly Asp
 325 330 335
 Val Leu Ser Glu Ala Gln Ile Lys Glu Cys Gln Glu Leu Gly Val Leu
 340 345 350
 Val Asp Arg Asp Asp Gln Gly Val Leu Leu Gln Ile Phe Thr Lys Pro
 355 360 365
 Val Gly Asp Arg Pro Thr Leu Phe Leu Glu Met Ile Gln Arg Ile Gly
 370 375 380
 Cys Met Glu Lys Asp Glu Arg Gly Glu Glu Tyr Gln Lys Gly Gly Cys
 385 390 395 400
 Gly Gly Phe Gly Lys Gly Asn Phe Ser Glu Leu Phe Lys Ser Ile Glu
 405 410 415
 Asp Tyr Glu Lys Ser Leu Glu Ala Lys Gln Ser Ala Ala Val Gln Gly
 420 425 430

Ser

<210> 39
 <211> 317
 <212> PRT
 <213> Synechocystis sp.

<400> 39
 Met Val Tyr His Val Arg Pro Lys His Ala Leu Phe Leu Ala Phe Tyr
 1 5 10 15

Cys Tyr Phe Ser Leu Leu Thr Met Ala Ser Ala Thr Ile Ala Ser Ala
 20 25 30
 Asp Leu Tyr Glu Lys Ile Lys Asn Phe Tyr Asp Asp Ser Ser Gly Leu
 35 40 45
 Trp Glu Asp Val Trp Gly Glu His Met His His Gly Tyr Tyr Gly Pro
 50 55 60
 His Gly Thr Tyr Arg Ile Asp Arg Arg Gln Ala Gln Ile Asp Leu Ile
 65 70 75 80
 Lys Glu Leu Leu Ala Trp Ala Val Pro Gln Asn Ser Ala Lys Pro Arg
 85 90 95
 Lys Ile Leu Asp Leu Gly Cys Gly Ile Gly Gly Ser Ser Leu Tyr Leu
 100 105 110
 Ala Gln Gln His Gln Ala Glu Val Met Gly Ala Ser Leu Ser Pro Val
 115 120 125
 Gln Val Glu Arg Ala Gly Glu Arg Ala Arg Ala Leu Gly Leu Gly Ser
 130 135 140
 Thr Cys Gln Phe Gln Val Ala Asn Ala Leu Asp Leu Pro Phe Ala Ser
 145 150 155 160
 Asp Ser Phe Asp Trp Val Trp Ser Leu Glu Ser Gly Glu His Met Pro
 165 170 175
 Asn Lys Ala Gln Phe Leu Gln Glu Ala Trp Arg Val Leu Lys Pro Gly
 180 185 190
 Gly Arg Leu Ile Leu Ala Thr Trp Cys His Arg Pro Ile Asp Pro Gly
 195 200 205
 Asn Gly Pro Leu Thr Ala Asp Glu Arg Arg His Leu Gln Ala Ile Tyr
 210 215 220
 Asp Val Tyr Cys Leu Pro Tyr Val Val Ser Leu Pro Asp Tyr Glu Ala
 225 230 235 240
 Ile Ala Arg Glu Cys Gly Phe Gly Glu Ile Lys Thr Ala Asp Trp Ser
 245 250 255
 Val Ala Val Ala Pro Phe Trp Asp Arg Val Ile Glu Ser Ala Phe Asp
 260 265 270
 Pro Arg Val Leu Trp Ala Leu Gly Gln Ala Gly Pro Lys Ile Ile Asn
 275 280 285
 Ala Ala Leu Cys Leu Arg Leu Met Lys Trp Gly Tyr Glu Arg Gly Leu
 290 295 300
 Val Arg Phe Gly Leu Leu Thr Gly Ile Lys Pro Leu Val
 305 310 315
 <210> 40
 <211> 348

<212> PRT

<213> Arabidopsis thaliana

<400> 40

Met Lys Ala Thr Leu Ala Ala Pro Ser Ser Leu Thr Ser Leu Pro Tyr
 1 5 10 15

Arg Thr Asn Ser Ser Phe Gly Ser Lys Ser Ser Leu Leu Phe Arg Ser
 20 25 30

Pro Ser Ser Ser Ser Ser Val Ser Met Thr Thr Thr Arg Gly Asn Val
 35 40 45

Ala Val Ala Ala Ala Ala Thr Ser Thr Glu Ala Leu Arg Lys Gly Ile
 50 55 60

Ala Glu Phe Tyr Asn Glu Thr Ser Gly Leu Trp Glu Glu Ile Trp Gly
 65 70 75 80

Asp His Met His His Gly Phe Tyr Asp Pro Asp Ser Ser Val Gln Leu
 85 90 95

Ser Asp Ser Gly His Lys Glu Ala Gln Ile Arg Met Ile Glu Glu Ser
 100 105 110

Leu Arg Phe Ala Gly Val Thr Asp Glu Glu Glu Glu Lys Lys Ile Lys
 115 120 125

Lys Val Val Asp Val Gly Cys Gly Ile Gly Gly Ser Ser Arg Tyr Leu
 130 135 140

Ala Ser Lys Phe Gly Ala Glu Cys Ile Gly Ile Thr Leu Ser Pro Val
 145 150 155 160

Gln Ala Lys Arg Ala Asn Asp Leu Ala Ala Ala Gln Ser Leu Ser His
 165 170 175

Lys Ala Ser Phe Gln Val Ala Asp Ala Leu Asp Gln Pro Phe Glu Asp
 180 185 190

Gly Lys Phe Asp Leu Val Trp Ser Met Glu Ser Gly Glu His Met Pro
 195 200 205

Asp Lys Ala Lys Phe Val Lys Glu Leu Val Arg Val Ala Ala Pro Gly
 210 215 220

Gly Arg Ile Ile Ile Val Thr Trp Cys His Arg Asn Leu Ser Ala Gly
 225 230 235 240

Glu Glu Ala Leu Gln Pro Trp Glu Gln Asn Ile Leu Asp Lys Ile Cys
 245 250 255

Lys Thr Phe Tyr Leu Pro Ala Trp Cys Ser Thr Asp Asp Tyr Val Asn
 260 265 270

Leu Leu Gln Ser His Ser Leu Gln Asp Ile Lys Cys Ala Asp Trp Ser
 275 280 285

Glu Asn Val Ala Pro Phe Trp Pro Ala Val Ile Arg Thr Ala Leu Thr
 290 295 300

Trp Lys Gly Leu Val Ser Leu Leu Arg Ser Gly Met Lys Ser Ile Lys
305 310 315 320

Gly Ala Leu Thr Met Pro Leu Met Ile Glu Gly Tyr Lys Lys Gly Val
325 330 335

Ile Lys Phe Gly Ile Ile Thr Cys Gln Lys Pro Leu
340 345

<210> 41

<211> 434

<212> PRT

<213> Hordeum vulgare

<400> 41

Met Pro Pro Thr Pro Thr Thr Pro Ala Ala Thr Gly Ala Ala Ala Ala
1 5 10 15

Val Thr Pro Glu His Ala Arg Pro His Arg Met Val Arg Phe Asn Pro
20 25 30

Arg Ser Asp Arg Phe His Thr Leu Ser Phe His His Val Glu Phe Trp
35 40 45

Cys Ala Asp Ala Ala Ser Ala Ala Gly Arg Phe Ala Phe Ala Leu Gly
50 55 60

Ala Pro Leu Ala Ala Arg Ser Asp Leu Ser Thr Gly Asn Ser Ala His
65 70 75 80

Ala Ser Gln Leu Leu Arg Ser Gly Ser Leu Ala Phe Leu Phe Thr Ala
85 90 95

Pro Tyr Ala Asn Gly Cys Asp Ala Ala Thr Ala Ser Leu Pro Ser Phe
100 105 110

Ser Ala Asp Ala Ala Arg Arg Phe Ser Ala Asp His Gly Ile Ala Val
115 120 125

Arg Ser Val Ala Leu Arg Val Ala Asp Ala Ala Glu Ala Phe Arg Ala
130 135 140

Ser Arg Arg Arg Gly Ala Arg Pro Ala Phe Ala Pro Val Asp Leu Gly
145 150 155 160

Arg Gly Phe Ala Phe Ala Glu Val Glu Leu Tyr Gly Asp Val Val Leu
165 170 175

Arg Phe Val Ser His Pro Asp Gly Thr Asp Val Pro Phe Leu Pro Gly
180 185 190

Phe Glu Gly Val Thr Asn Pro Asp Ala Val Asp Tyr Gly Leu Thr Arg
195 200 205

Phe Asp His Val Val Gly Asn Val Pro Glu Leu Ala Pro Ala Ala Ala
210 215 220

Tyr Ile Ala Gly Phe Thr Gly Phe His Glu Phe Ala Glu Phe Thr Ala
225 230 235 240

Glu Asp Val Gly Thr Thr Glu Ser Gly Leu Asn Ser Val Val Leu Ala
 245 250 255
 Asn Asn Ser Glu Gly Val Leu Leu Pro Leu Asn Glu Pro Val His Gly
 260 265 270
 Thr Lys Arg Arg Ser Gln Ile Gln Thr Phe Leu Glu His His Gly Gly
 275 280 285
 Pro Gly Val Gln His Ile Ala Val Ala Ser Ser Asp Val Leu Arg Thr
 290 295 300
 Leu Arg Lys Met Arg Ala Arg Ser Ala Met Gly Gly Phe Asp Phe Leu
 305 310 315 320
 Pro Pro Pro Leu Pro Lys Tyr Tyr Glu Gly Val Arg Arg Leu Ala Gly
 325 330 335
 Asp Val Leu Ser Glu Ala Gln Ile Lys Glu Cys Gln Glu Leu Gly Val
 340 345 350
 Leu Val Asp Arg Asp Asp Gln Gly Val Leu Leu Gln Ile Phe Thr Lys
 355 360 365
 Pro Val Gly Asp Arg Pro Thr Leu Phe Leu Glu Met Ile Gln Arg Ile
 370 375 380
 Gly Cys Met Glu Lys Asp Glu Arg Gly Glu Glu Tyr Gln Lys Gly Gly
 385 390 395 400
 Cys Gly Gly Phe Gly Lys Gly Asn Phe Ser Glu Leu Phe Lys Ser Ile
 405 410 415
 Glu Asp Tyr Glu Lys Ser Leu Glu Ala Lys Gln Ser Ala Ala Val Gln
 420 425 430

Gly Ser

<210> 42
 <211> 442
 <212> PRT
 <213> Daucus carota

<400> 42
 Met Gly Lys Lys Gln Ser Glu Ala Glu Ile Leu Ser Ser Asn Ser Ser
 1 5 10 15
 Asn Thr Ser Pro Ala Thr Phe Lys Leu Val Gly Phe Asn Asn Phe Val
 20 25 30
 Arg Ala Asn Pro Lys Ser Asp His Phe Ala Val Lys Arg Phe His His
 35 40 45
 Ile Glu Phe Trp Cys Gly Asp Ala Thr Asn Thr Ser Arg Arg Phe Ser
 50 55 60
 Trp Gly Leu Gly Met Pro Leu Val Ala Lys Ser Asp Leu Ser Thr Gly
 65 70 75 80

Asn Ser Val His Ala Ser Tyr Leu Val Arg Ser Ala Asn Leu Ser Phe
 85 90 95
 Val Phe Thr Ala Pro Tyr Ser Pro Ser Thr Thr Thr Ser Ser Gly Ser
 100 105 110
 Ala Ala Ile Pro Ser Phe Ser Ala Ser Gly Phe His Ser Phe Ala Ala
 115 120 125
 Lys His Gly Leu Ala Val Arg Ala Ile Ala Leu Glu Val Ala Asp Val
 130 135 140
 Ala Ala Ala Phe Glu Ala Ser Val Ala Arg Gly Ala Arg Pro Ala Ser
 145 150 155 160
 Ala Pro Val Glu Leu Asp Asp Gln Ala Trp Leu Ala Glu Val Glu Leu
 165 170 175
 Tyr Gly Asp Val Val Leu Arg Phe Val Ser Phe Gly Arg Glu Glu Gly
 180 185 190
 Leu Phe Leu Pro Gly Phe Glu Ala Val Glu Gly Thr Ala Ser Phe Pro
 195 200 205
 Asp Leu Asp Tyr Gly Ile Arg Arg Leu Asp His Ala Val Gly Asn Val
 210 215 220
 Thr Glu Leu Gly Pro Val Val Glu Tyr Ile Lys Gly Phe Thr Gly Phe
 225 230 235 240
 His Glu Phe Ala Glu Phe Thr Ala Glu Asp Val Gly Thr Leu Glu Ser
 245 250 255
 Gly Leu Asn Ser Val Val Leu Ala Asn Asn Glu Glu Met Val Leu Leu
 260 265 270
 Pro Leu Asn Glu Pro Val Tyr Gly Thr Lys Arg Lys Ser Gln Ile Gln
 275 280 285
 Thr Tyr Leu Glu His Asn Glu Gly Ala Gly Val Gln His Leu Ala Leu
 290 295 300
 Val Ser Glu Asp Ile Phe Arg Thr Leu Arg Glu Met Arg Lys Arg Ser
 305 310 315 320
 Cys Leu Gly Gly Phe Glu Phe Met Pro Ser Pro Pro Pro Thr Tyr Tyr
 325 330 335
 Lys Asn Leu Lys Asn Arg Val Gly Asp Val Leu Ser Asp Glu Gln Ile
 340 345 350
 Lys Glu Cys Glu Asp Leu Gly Ile Leu Val Asp Arg Asp Asp Gln Gly
 355 360 365
 Thr Leu Leu Gln Ile Phe Thr Lys Pro Val Gly Asp Arg Pro Thr Leu
 370 375 380
 Phe Ile Glu Ile Ile Gln Arg Val Gly Cys Met Leu Lys Asp Asp Ala
 385 390 395 400

33

Val Ala Gly Phe Thr Gly Phe His Gln Phe Ala Glu Phe Thr Ala Asp
 245 250 255
 Asp Val Gly Thr Ala Glu Ser Gly Leu Asn Ser Ala Val Leu Ala Ser
 260 265 270
 Asn Asp Glu Met Val Leu Leu Pro Ile Asn Glu Pro Val His Gly Thr
 275 280 285
 Lys Arg Lys Ser Gln Ile Gln Thr Tyr Leu Glu His Asn Glu Gly Ala
 290 295 300
 Gly Leu Gln His Leu Ala Leu Met Ser Glu Asp Ile Phe Arg Thr Leu
 305 310 315 320
 Arg Glu Met Arg Lys Arg Ser Ser Ile Gly Gly Phe Asp Phe Met Pro
 325 330 335
 Ser Pro Pro Pro Thr Tyr Tyr Gln Asn Leu Lys Lys Arg Val Gly Asp
 340 345 350
 Val Leu Ser Asp Asp Gln Ile Lys Glu Cys Glu Glu Leu Gly Ile Leu
 355 360 365
 Val Asp Arg Asp Asp Gln Gly Thr Leu Leu Gln Ile Phe Thr Lys Pro
 370 375 380
 Leu Gly Asp Arg Pro Thr Ile Phe Ile Glu Ile Ile Gln Arg Val Gly
 385 390 395 400
 Cys Met Met Lys Asp Glu Glu Gly Lys Ala Tyr Gln Ser Gly Gly Cys
 405 410 415
 Gly Gly Phe Gly Lys Gly Asn Phe Ser Glu Leu Phe Lys Ser Ile Glu
 420 425 430
 Glu Tyr Glu Lys Thr Leu Glu Ala Lys Gln Leu Val Gly
 435 440 445

```

SEC 10 MO:26 0MSNPATPPIVAIKELTKEGLTSLARSQVETVGGANVPLVIEGVEKELKFPITTCB
          301
SEC 10 MO:39 KPLN-----
SEC 10 MO:40 KPL-----
SEC 10 MO:38 KFORAIEGEGMAASPYE.-
SEC 10 MO:28 K-----PE-----
SEC 10 MO:08 SYVYDTYFKTD.-
SEC 10 MO:22 KPGAA--NA---KAALALIC
SEC 10 MO:24

```

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

INTERNATIONAL SEARCH REPORT

Internat : Application No

PCT/US 99/28588

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C12N9/10 C12N9/02 C12N15/82 C12N15/29 C12N15/63
C12N15/83 C12Q1/48 C12Q1/26 C12Q1/68 G01N33/50

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C12N C12Q

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>DATABASE EMBL [Online] Accession No. D64004, 4 October 1995 (1995-10-04) "Synechocystis sp. PCC6803 complete genome, delta(24)-sterol C-methyltransferase" XP002135563 see p. 22, l. 38 of present description: 1001725 corresponds to BAA10562 and D64004 the whole document</p> <p style="text-align: center;">--- -/--</p>	1,3-11, 13-25

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"Z" document member of the same patent family

Date of the actual completion of the international search

12 April 2000

Date of mailing of the international search report

101 09 00

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Herrmann, K

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 99/28588

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	BARTLEY, G.E. ET AL.: "Cloning of an Arabidopsis thaliana cDNA for p-hydroxyphenylpyruvate dioxygenase (Accession No. U89267)" PLANT PHYSIOL., vol. 113, 1997, pages 1465-1465, XP002135561 the whole document	18
A	& DATABASE EMBL [Online] Accession No. U89267, 19 March 1997 (1997-03-19) "Arabidopsis thaliana p-hydroxyphenylpyruvate dioxygenase (HPD) mRNA, complete cds" the whole document	
A	SHINTANI, D. AND DELLAPENNA, D.: "A new paradigm for plant biochemistry: Nutritional Genomics. Vitamin E synthesis and human nutrition as an example.", Abstract of the oral presentation at the annual meeting of The American Society of Plant Physiology, 28.06.1998 XP002135562 the whole document	
P,X	SHINTANI, D. AND DELLAPENNA, D.: "Elevating the vitamin E content of plants through metabolic engineering" SCIENCE, vol. 282, no. 5396, 11 December 1998 (1998-12-11), pages 2098-2100, XP000887122 the whole document	
P,A	WO 99 04622 A (UNIV NEVADA) 4 February 1999 (1999-02-04)	

INTERNATIONAL SEARCH REPORT

Inter. application No.
PCT/US 99/28588

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:

3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

See additional sheet

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.

2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.

3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:

4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

1, 3-11, 13-25 (all partially)

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

1. Claims: Invention 1: Claims 1, 3-11, 13-25 (all partially)

Corn gamma-tocopherol methyltransferase:

Polynucleotide encoding a polypeptide as in SEQ ID NO: 2 or 22, polypeptide as in SEQ ID NO:2 or 22, polynucleotide comprising a nucleotide sequence as in SEQ ID NO:1 or 21. Subject-matter relating to corn gamma-tocopherol methyltransferase.

2. Claims: Invention 2: Claims 1, 3-11, 13-25 (all partially)

Rice gamma-tocopherol methyltransferase:

Polynucleotide encoding a polypeptide as in SEQ ID NO:4, 6, 24 or 26, polypeptide as in SEQ ID NO:4, 6, 24 or 26, polynucleotide comprising a nucleotide sequence as in SEQ ID NO:3, 5, 23 or 25. Subject-matter relating to rice gamma-tocopherol methyltransferase.

3. Claims: Invention 3: Claims 1, 3-11, 13-25 (all partially)

Soybean gamma-tocopherol methyltransferase:

Polynucleotide encoding a polypeptide as in SEQ ID NO:8 or 28, polypeptide as in SEQ ID NO:8 or 28, polynucleotide comprising a nucleotide sequence as in SEQ ID NO:7 or 27. Subject-matter relating to soybean gamma-tocopherol methyltransferase.

4. Claims: Invention 4: Claims 1, 3-11, 13-25 (all partially)

Wheat gamma-tocopherol methyltransferase:

Polynucleotide encoding a polypeptide as in SEQ ID NO:8 or 28, polypeptide as in SEQ ID NO:10 or 30, polynucleotide comprising a nucleotide sequence as in SEQ ID NO:10 or 30. Subject-matter relating to wheat gamma-tocopherol methyltransferase.

5. Claims: Invention 5: Claims 2-10, 12-19, 21-25 (all partially)

Rice 4-hydroxyphenylpyruvate dioxygenase:

Polynucleotide encoding a polypeptide as in SEQ ID NO:12, 14 or 34, polypeptide as in SEQ ID NO:12, 14 or 34, polynucleotide comprising a nucleotide sequence as in SEQ ID NO:11, 13 or 33. Subject-matter relating to rice 4-hydroxyphenylpyruvate dioxygenase.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

6. Claims: Invention 6: Claims 2-10, 12-19,
21-25 (all partially)

Soybean 4-hydroxyphenylpyruvate dioxygenase:

Polynucleotide encoding a polypeptide as in SEQ ID NO:16 or 36, polypeptide as in SEQ ID NO:16 or 36, polynucleotide comprising a nucleotide sequence as in SEQ ID NO:15 or 35. Subject-matter relating to soybean 4-hydroxyphenylpyruvate dioxygenase.

7. Claims: Invention 7: Claims 2-10, 12-19,
21-25 (all partially)

Veronica 4-hydroxyphenylpyruvate dioxygenase:

Polynucleotide encoding a polypeptide as in SEQ ID NO:18, polypeptide as in SEQ ID NO:18, polynucleotide comprising a nucleotide sequence as in SEQ ID NO:17. Subject-matter relating to veronica 4-hydroxyphenylpyruvate dioxygenase.

8. Claims: Invention 8: Claims 2-10, 12-19,
21-25 (all partially)

Wheat 4-hydroxyphenylpyruvate dioxygenase:

Polynucleotide encoding a polypeptide as in SEQ ID NO:20 or 38, polypeptide as in SEQ ID NO:20 or 38, polynucleotide comprising a nucleotide sequence as in SEQ ID NO:19 or 37. Subject-matter relating to wheat 4-hydroxyphenylpyruvate dioxygenase.

9. Claims: Invention 9: Claims 2-10, 12-19,
21-25 (all partially)

Catalpa 4-hydroxyphenylpyruvate dioxygenase:

Polynucleotide encoding a polypeptide as in SEQ ID NO:32, polypeptide as in SEQ ID NO:32, polynucleotide comprising a nucleotide sequence as in SEQ ID NO:31. Subject-matter relating to catalpa 4-hydroxyphenylpyruvate dioxygenase.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 99/28588

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9904622 A	04-02-1999	AU 8506198 A EP 1009812 A	16-02-1999 21-06-2000
